



TECHNISCHE UNIVERSITÄT CHEMNITZ

Sonderforschungsbereich 393

Parallele Numerische Simulation für Physik und Kontinuumsmechanik

Cornelia Pester

Hamiltonian eigenvalue symmetry for quadratic operator eigenvalue problems

Preprint SFB393/04-09

Abstract When the eigenvalues of a given eigenvalue problems are symmetric with respect to the real and the imaginary axes, we speak about a Hamiltonian eigenvalue symmetry or a Hamiltonian structure of the spectrum. This property can be exploited for an efficient computation of the eigenvalues. For some elliptic boundary value problems it is known that the derived eigenvalue problems have this Hamiltonian symmetry. Without having a specific application in mind, we trace the question, under which assumptions the spectrum of a given quadratic eigenvalue problem possesses the Hamiltonian structure.

Key Words nonlinear eigenvalue problems, Hamiltonian eigenvalue symmetry, spectral theory, operator pencil

AMS subject classification 35P30, 37K05, 65F15

Preprintreihe des Chemnitzer SFB 393

ISSN 1619-7178 (Print)

ISSN 1619-7186 (Internet)

SFB393/04-09

October 2004

Contents

1	Introduction	1
2	Operators in Hilbert spaces	3
2.1	General notation	3
2.2	Compact operators and Fredholm operators	4
3	The quadratic eigenvalue problem	5
3.1	Definitions and assumptions	5
3.2	The eigenvalue problem for an operator pencil	7
3.3	Compactness of \mathcal{M} and \mathcal{G}	9
3.4	Proof: $\lambda \in \sigma(\mathcal{B}) \implies -\bar{\lambda} \in \sigma(\mathcal{B})$	9
3.5	Proof: $\lambda \in \sigma(\mathcal{B}) \implies \bar{\lambda} \in \sigma(\mathcal{B})$	10
3.6	The Hamiltonian eigenvalue symmetry	11
4	Examples	11
5	Summary	12

Author's address:

Cornelia Pester
Institut für Mathematik
Fakultät für Bauingenieur- und Vermessungswesen
Universität der Bundeswehr München
85577 Neubiberg, Germany

cornelia.pesther@unibw-muenchen.de
<http://www.bauv.unibw-muenchen.de/pester/>

1 Introduction

Non-linear eigenvalue problems arise in several fields of mathematics including the analysis and numerical solution of higher order systems of ordinary differential equations, for example, in the robot model with electric motors in the joints, see Schiehlen (1990) or Mehrmann and Watkins (2002). Although solvers for quadratic eigenvalue problems are known, see, for example, Bai (2002), the common way is to linearize the eigenvalue problem and to transform it to a standard eigenvalue problem, for whose efficient solution many algorithms exist; a good summary is given by Watkins (2002).

We focus on polynomial eigenvalue problems of order two. Quadratic eigenvalue problems are treated, for instance, in the analysis and algebra communities as well as in the engineering community. Depending on the underlying model problem, the resulting quadratic operator eigenvalue problem might have a considerable structure which can be exploited in the further analysis and computations. Our numerical interest in a detailed analysis of such a structure arose from papers including work by Leguillon (1995), Kozlov, Maz'ya, and Roßmann (2000), Apel, Sändig, and Solov'ev (2002b) and Apel, Mehrmann, and Watkins (2002a), who studied the linear elasticity problem in the neighbourhood of polyhedral corners. The computation of the singular part of the solution to the linear elasticity problem near a polyhedral corner involves the solution of a quadratic eigenvalue problem of the form: Find $\lambda \in \mathbb{C}$, $u \in \mathcal{H}$, such that for all $v \in \mathcal{H}$

$$\lambda^2 m(u, v) + \lambda g(u, v) = k(u, v), \quad (1)$$

where \mathcal{H} is a given Hilbert space. The functions m , g , k are sesquilinear forms defined on $\mathcal{H} \times \mathcal{H}$, where m and k are Hermitian and g is skew-Hermitian.

It is well-known that such eigenvalue problems which are associated with elliptic boundary value problems have a considerable structure: the eigenvalues appear in pairs $(\lambda, -\lambda)$ if λ is real or purely imaginary and in quadruplets $(\lambda, -\lambda, \bar{\lambda}, -\bar{\lambda})$ otherwise, see Figure 1. This property is called Hamiltonian eigenvalue symmetry or Hamiltonian structure of the spectrum, motivated by the spectral properties of a Hamiltonian matrix, see, for example, Mehrmann and Watkins (2002).




Figure 1: The Hamiltonian structure of eigenvalues

The Hamiltonian structure for eigenvalue problems which are associated with certain elliptic boundary value problems in the neighbourhood of three-dimensional polyhedral corners was observed, for instance, by Leguillon (1995) as a generalization of the two-dimensional case or by Kozlov, Maz'ya, and Roßmann (2000) (see also referenced therein) who applied a Mellin transformation to the given boundary value problem.

Under which assumptions is the Hamiltonian eigenvalue symmetry guaranteed for problem (1), in general? The focus in this paper lies on eigenvalue problems of type (1) without any specific application in the background. Obviously, the Hamiltonian structure is not always given. Consider, for instance, $g(u, v) = i m(u, v)$, $k(u, v) \equiv 0$, where $i \in \mathbb{C}$ with $i^2 = -1$. Then $\lambda = -i$ is an eigenvalue of the problem

$$\lambda^2 m(u, v) + \lambda i m(u, v) = 0,$$

but $\lambda = i$ is not. We traced the spectral properties of problem (1) and formulated a series of conditions which imply the Hamiltonian structure. Since the conditions are formulated in an easy-to-check state, they are sufficient, but not urgently necessary as is demonstrated in some examples in Section 4. Still, they are satisfied in the practical applications that we have in mind.

The Hamiltonian eigenvalue symmetry is of interest in the further analysis of the problem like an efficient solution of its discrete formulation. Adapted Arnoldi and Lanczos algorithms can be applied then which exploit the underlying structure. For a description of such algorithms, we refer to Freund (1994); Benner and Faßbender (1997, 2000); Mehrmann and Watkins (2001); Apel, Mehrmann, and Watkins (2002a); Watkins (2004).

We could not find a self-contained proof or a summary of conditions for the Hamiltonian structure of the spectrum of a quadratic eigenvalue problem in the linear algebra, functional analysis or numerical analysis literature. On few pages, we give in this paper an overview of the most important ingredients and definitions, followed by a list of assumptions and a read-on-proof of the Hamiltonian eigenvalue symmetry, including a discussion of the necessity of the assumptions. Of course, there exist other methods to prove this result, suggested, for example, by Kozlov, Maz'ya, and Roßmann (2000), but they usually employ more background information of the given problem and refer to further literature.

As the derivation requires ingredients from several fields of the mathematical sciences, we give an overview of the most important terms in Section 2. For details, we refer to standard work in functional analysis, operator theory, spectral theory, algebra and numerical analysis. In Section 3, we discuss spectral properties of problem (1) and an associated eigenvalue problem for an operator pencil. We conclude the paper with a few examples including an application of the results in Section 4.

2 Operators in Hilbert spaces

2.1 General notation

Let H_R be a (separable) *real* Hilbert space and denote by H_C its complexification

$$H_C = H_R \times H_R = \{u = (u_1, u_2) \mid u_1, u_2 \in H_R\}$$

with the operations

$$\begin{aligned} (u_1, u_2) + (v_1, v_2) &= (u_1 + v_1, u_2 + v_2) \quad \forall u_i, v_i \in H_R, \\ (a + ib)(u, v) &= (au - bv, bu + av) \quad \forall a, b \in \mathbb{R}, \forall u, v \in H_R. \end{aligned}$$

Each element $u_R \in H_R$ shall be identified with the element $(u_R, 0) \in H_C$, so that $H_R \subset H_C$ and each element $u = (u_1, u_2) \in H_C$ can be written as $u = (u_1, 0) + i(u_2, 0) = u_1 + iu_2$. Moreover, we define the conjugate of $u = (u_1, u_2) \in H_C$ by

$$\bar{u} = \overline{u_1 + iu_2} = u_1 - iu_2.$$

Definition 2.1 (Sesquilinear form). A map $a : H_C \times H_C \rightarrow \mathbb{C}$ is called a sesquilinear form over H_C , if

$$a(\mu u, v) = \mu a(u, v), \quad a(u, \mu v) = \bar{\mu} a(u, v) \quad \forall \mu \in \mathbb{C}, \forall u, v \in H_C$$

and

$$\begin{aligned} a(u + w, v) &= a(u, v) + a(w, v) \quad \forall u, v, w \in H_C \\ a(u, v + w) &= a(u, v) + a(u, w) \quad \forall u, v, w \in H_C. \end{aligned}$$

Let $\langle \cdot, \cdot \rangle_{H_C} : H_C \times H_C \rightarrow \mathbb{C}$ denote an inner product on H_C , that is a Hermitian, positive definite sesquilinear form. A *norm* in H_C is then given by $\|u\|_{H_C} := \sqrt{\langle u, u \rangle_{H_C}}$.

Definition 2.2 (Real operator). A linear, bounded operator $\mathcal{A} : H_C \rightarrow H_C$ is called a real operator, if $\mathcal{A}(H_R) \subset H_R$, i.e., if $\mathcal{A}u_R \in H_R$ for all $u_R \in H_R$.

Definition 2.3 (Spectrum of an operator pencil). Let $\mathcal{A} : \mathbb{C} \rightarrow \mathcal{L}(H_C, H_C)$ be an operator pencil; that is, $\mathcal{A}(\alpha) : H_C \rightarrow H_C$ for fixed $\alpha \in \mathbb{C}$. The set

$$\Sigma(\mathcal{A}) := \{\alpha \in \mathbb{C} \mid \mathcal{A}(\alpha) \text{ has no bounded inverse}\}$$

is called the spectrum of $\mathcal{A}(\cdot)$. The number $\alpha_0 \in \Sigma(\mathcal{A})$ is called eigenvalue of $\mathcal{A}(\cdot)$, if there exists an element $u_{\alpha_0} \in H_C \setminus \{0\}$ so that $\mathcal{A}(\alpha_0)u_{\alpha_0} = 0$. We denote the set of all eigenvalues of $\mathcal{A}(\cdot)$ by $\sigma(\mathcal{A})$:

$$\sigma(\mathcal{A}) := \{\alpha \in \mathbb{C} \mid \exists u \in H_C \setminus \{0\}, \text{ so that } \mathcal{A}(\alpha)u = 0\}.$$

Eigenvalue problems for operator pencils have already been considered, for example, by Kozlov, Maz'ya, and Roßmann (2000).

Remark 2.4. We say that an operator $\mathcal{J} : X \rightarrow Y$ is invertible (or that \mathcal{J}^{-1} exists), if for all $y \in Y$ there is a unique element $x \in X$, so that $\mathcal{J}x = y$. In some books, the invertibility is defined by substituting Y by the range $R(\mathcal{J})$ of \mathcal{J} ; then, the operator \mathcal{J} is surjective by definition and the invertibility is equivalent to the injectivity of \mathcal{J} .

Note that the sets $\Sigma(\mathcal{A})$ and $\sigma(\mathcal{A})$ are equal for finite-dimensional operators; but they might differ for operators in infinite-dimensional spaces. The set $\Sigma(\mathcal{A}) \setminus \sigma(\mathcal{A})$ is called the continuous spectrum of $\mathcal{A}(\cdot)$. The equivalence of $\ker(\mathcal{A}(\alpha)) = \{0\}$ and the invertibility of $\mathcal{A}(\alpha)$ is given only for special linear operators $\mathcal{A}(\cdot)$, see Section 2.2.

2.2 Compact operators and Fredholm operators

In this subsection, we give the definitions and, for our purposes, the most important properties of compact and Fredholm operators. For details and proofs, consult Alt (1999); Werner (1997); Prößdorf (1974); Cotlar and Cignoli (1974) or any standard book on functional analysis or operator theory.

Let H be a separable Hilbert space with an inner product $\langle \cdot, \cdot \rangle_H : H \times H \rightarrow \mathbb{C}$ on H .

Definition 2.5 (Compact operators). A linear, bounded operator $\mathcal{C} : H \rightarrow H$ is called compact, if for each bounded sequence $\{x_n\}$ in H , there is a subsequence $\{x_{n_k}\}$ so that $\mathcal{C}x_{n_k}$ converges in H , that is, if

$$\|x_n\|_H < \infty \quad \forall n \quad \implies \quad \exists x_{n_k}, y^* \in H : \|\mathcal{C}x_{n_k} - y^*\|_H \longrightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Remark 2.6. Let $K(H)$ denote the set of all compact operators on H . Then $K(H)$ is a subspace of the space $\mathcal{L}(H)$ of all bounded, linear operators on H , see Werner (1997, Satz II.3.2 (a)). This means, in particular, that the sum of two compact operators as well as a scalar multiple of a compact operator is in $K(H)$ and therefore compact.

Definition 2.7 (Compact embedding). We say that a space V is compactly embedded into a space $H \supset V$, if the corresponding embedding operator is compact, that is, if for each bounded sequence $\{u_n\}$ in V there is a subsequence $\{u_{n_k}\}$ with strong convergence in H .

We denote by $R(\mathcal{A})$ and $\ker(\mathcal{A})$ the range and the kernel of the linear, bounded operator $\mathcal{A} : H \rightarrow H$, respectively, and by $R(\mathcal{A}^*)$ and $\ker(\mathcal{A}^*)$ the range and the kernel of its adjoint.

Lemma 2.8. The relations

$$\ker(\mathcal{A}) \oplus \overline{R(\mathcal{A}^*)} = H \quad \text{and} \quad \ker(\mathcal{A}^*) \oplus \overline{R(\mathcal{A})} = H$$

hold for each linear, bounded operator $\mathcal{A} : H \rightarrow H$.

Lemma 2.8 is a well-known result in functional analysis, see, for example, Werner (1997, Satz III.4.5).

Definition 2.9 (Fredholm operators). A linear, bounded operator \mathcal{T} is called Fredholm operator if $\dim \ker \mathcal{T} < \infty$ and $\dim \ker \mathcal{T}^* < \infty$. For a Fredholm operator \mathcal{T} , the number $\text{Ind } \mathcal{T} = \dim \ker \mathcal{T} - \dim \ker \mathcal{T}^*$ is called the index of \mathcal{T} .

Lemma 2.10. For each linear, bounded operator \mathcal{T} the following properties are equivalent:

- (a) \mathcal{T} is Fredholm and $\text{Ind } \mathcal{T} = 0$.
- (b) There is a compact operator \mathcal{C} and an invertible operator \mathcal{J} , so that $\mathcal{T} = \mathcal{J} - \mathcal{C}$.

Remark 2.11. Suppose that (a) is true in Lemma 2.10. Then $\dim \ker \mathcal{T} = \dim \ker \mathcal{T}^*$ and thus $\overline{\ker \mathcal{T}} = \{0\}$ if and only if $\overline{\ker \mathcal{T}^*} = \{0\}$. As a consequence of Lemma 2.8, we have that $R(\mathcal{T}) = H$ if and only if $R(\mathcal{T}^*) = H$. Hence, \mathcal{T} is invertible if and only if $\ker \mathcal{T} = \{0\}$. Moreover, \mathcal{T} is invertible if and only if \mathcal{T}^* is invertible.

3 The quadratic eigenvalue problem

3.1 Definitions and assumptions

Let $m, g, k : H_C \times H_C \rightarrow \mathbb{C}$ be sesquilinear forms over the Hilbert space H_C with

$$k(u, v) = \overline{k(v, u)}, \quad g(u, v) = -\overline{g(v, u)}, \quad m(u, v) = \overline{m(v, u)} \quad \forall u, v \in H_C.$$

We consider the quadratic eigenvalue problem (1): Find $\lambda \in \mathbb{C}$, $u \in H_C \setminus \{0\}$, such that

$$\lambda^2 m(u, v) + \lambda g(u, v) = k(u, v) \quad \forall v \in H_C.$$

A solution $(\lambda, u) \in \mathbb{C} \times (H_C \setminus \{0\})$ to problem (1) is called an *eigenpair* of problem (1); the number λ is then called *eigenvalue* and u is called *eigenelement*.

Definition 3.1 (Hamiltonian eigenvalue symmetry). We say that problem (1) possesses a Hamiltonian eigenvalue symmetry or that its spectrum has a Hamiltonian structure, if the eigenvalues are placed symmetric with respect to the real and the imaginary axes; that is, if $\lambda \in \mathbb{C}$ is an eigenvalue of problem (1), then so are $-\lambda$, $\bar{\lambda}$, $-\bar{\lambda}$.

Remark 3.2. A real two-by-two block matrix H is called Hamiltonian, if

$$(JH)^\top = JH \quad \text{with} \quad J = \begin{pmatrix} O & I \\ -I & O \end{pmatrix}.$$

The term Hamiltonian eigenvalue symmetry is motivated by the structure of the spectrum of Hamiltonian matrices.

The question is under which conditions on m, g, k and H_C the Hamiltonian eigenvalue symmetry is given. In the derivation of the desired properties, we followed the ideas of Apel, Sändig, and Solov'ev (2002b), where a quadratic eigenvalue problem of type (1) is treated and transformed to an eigenvalue problem for an operator pencil.

In the following assumptions, we summarize conditions that are sufficient for the Hamiltonian structure. They are usually easy to check and satisfied in the applications that we have in mind.

(I) $V := H_C := H_R \times H_R$ is the complexification of a separable real Hilbert space H_R , and $H \supset V$ is a Hilbert space into which V is compactly embedded.

(II) The functions $m : H \times H \rightarrow \mathbb{C}$, $g, k : V \times V \rightarrow \mathbb{C}$ are sesquilinear forms with

$$m(u, v) = \overline{m(v, u)} \quad \forall u, v \in H, \quad k(u, v) = \overline{k(v, u)} \quad \forall u, v \in V,$$

(III) The functions m, g, k are real, that is,

$$m(\bar{u}, \bar{v}) = \overline{m(u, v)}, \quad g(\bar{u}, \bar{v}) = \overline{g(u, v)}, \quad k(\bar{u}, \bar{v}) = \overline{k(u, v)} \quad \forall u, v \in V.$$

(IV) The sesquilinear form m satisfies

$$|m(u, v)| \leq c \|u\|_H \|v\|_H \quad \forall u, v \in H.$$

(V) There is a sesquilinear form $d : H \times V \rightarrow \mathbb{C}$ so that $g(u, v) = \overline{d(v, u)} - d(u, v)$ for all $u, v \in V$ and

$$|d(u, v)| \leq c \|u\|_H \|v\|_V \quad \forall u, v \in V.$$

(VI) The operator $\mathcal{K} : V \rightarrow V$ induced by $k(u, v) = \langle \mathcal{K}u, v \rangle_V \quad \forall u, v \in V$ is invertible; and for each fixed element $u \in H$, there is a constant $c_k > 0$, so that $|k(u, v)| \leq c_k \|v\|_V$ for all $v \in V$.

Alternatively, it is sufficient that the sesquilinear form k is positive definite, that is,

$$k(u, u) \geq 0 \quad \forall u \in V, \quad k(u, u) = 0 \iff u = 0.$$

In the following, V and H denote the Hilbert spaces introduced by assumption (I).

Remark 3.3. *Given a skew-Hermitian sesquilinear form $g : V \times V \rightarrow \mathbb{C}$, there is always a map $d : V \times V \rightarrow \mathbb{C}$ so that $g(u, v) = \overline{d(v, u)} - d(u, v)$, for example, $d(u, v) = -\frac{1}{2}g(u, v)$. Assumption (V), however, demands that the sesquilinear form d is defined on a larger space, on $H \times V$, so that $d = -\frac{1}{2}g$ is not necessarily the desired function. That is why we cannot simply omit d or g . Still, the function g has to be defined only on the space $V \times V$, whereas m has the largest domain $H \times H$ due to assumption (II).*

The condition (III) that the functions m, g and k are real does not mean that they are real-valued, but only that they have real values for real arguments. This is equivalent to

$$m(u_R, v_R) \in \mathbb{R}, \quad g(u_R, v_R) \in \mathbb{R}, \quad k(u_R, v_R) \in \mathbb{R}, \quad \forall u_R, v_R \in H_R.$$

If $k(\cdot, \cdot)$ defines an inner product on V , that is, if $k(\cdot, \cdot)$ is positive definite, we can choose $\langle u, v \rangle_V = k(u, v)$ for all $u, v \in V$. Then, the operator \mathcal{K} in condition (VI) equals the identity operator \mathcal{I} on V and is therefore invertible.

For the inner products $\langle \cdot, \cdot \rangle_V$ in V and $\langle \cdot, \cdot \rangle_H$ in H , the corresponding norms are given by $\|u\|_V^2 = \langle u, u \rangle_V$ and $\|u\|_H^2 = \langle u, u \rangle_H$. Due to assumption (I), the space V is embedded into H . Thus, the relation

$$\|u\|_H \leq c\|u\|_V \quad (2)$$

holds for all $u \in V$, where c is a generic constant independent of u .

In the remaining section, we prove that the Hamiltonian eigenvalue symmetry is actually given under these assumptions. When a weaker condition is sufficient in one or another step, we will point this out.

3.2 The eigenvalue problem for an operator pencil

Corresponding to the sesquilinear forms m , g and k , we introduce the linear operators $\mathcal{M}, \mathcal{G}, \mathcal{K} : V \rightarrow V$ by

$$\begin{aligned} \mathcal{M}u \in V, \quad \langle \mathcal{M}u, v \rangle_V &= m(u, v) \quad \forall v \in V, \\ \mathcal{G}u \in V, \quad \langle \mathcal{G}u, v \rangle_V &= g(u, v) \quad \forall v \in V, \\ \mathcal{K}u \in V, \quad \langle \mathcal{K}u, v \rangle_V &= k(u, v) \quad \forall v \in V. \end{aligned}$$

The adjoint operators \mathcal{M}^* , \mathcal{G}^* and \mathcal{K}^* are given by

$$\begin{aligned} \langle \mathcal{M}u, v \rangle_V &= \langle u, \mathcal{M}^*v \rangle_V \quad \forall u, v \in V, \\ \langle \mathcal{G}u, v \rangle_V &= \langle u, \mathcal{G}^*v \rangle_V \quad \forall u, v \in V, \\ \langle \mathcal{K}u, v \rangle_V &= \langle u, \mathcal{K}^*v \rangle_V \quad \forall u, v \in V. \end{aligned}$$

Lemma 3.4. *The operators \mathcal{M} , \mathcal{G} and \mathcal{K} are well-defined.*

Proof. The Riesz representation theorem states: Let $\langle \cdot, \cdot \rangle_X$ be an inner product on a Hilbert space X . Then for each bounded linear functional $F \in X^*$, there is a unique element $u_F \in X$, so that $\langle v, u_F \rangle_X = F(v)$ for all $v \in X$, see, for example, Alt (1999).

We choose $X := V$ and $\langle \cdot, \cdot \rangle_X := \langle \cdot, \cdot \rangle_V$ as well as $F(v) := F_{m, u_0}(v) := m(u_0, v)$ for a fixed element $u_0 \in V$. Indeed, $F_{m, u_0}(v)$ is a linear functional; but for its boundedness, we have to assume that $|m(u_0, v)| \leq c\|v\|_V$ for all $v \in V$ and for each fixed $u_0 \in V$ with a generic constant c independent of v . Then, there is a unique element $u_{F_{m, u_0}} \in V$, so that $\langle v, u_{F_{m, u_0}} \rangle_V = \overline{m(u_0, v)}$, and therefore, $\langle u_{F_{m, u_0}}, v \rangle_V = m(u_0, v)$ for all $v \in V$. Consequently, $\mathcal{M}u_0$ is given by $\mathcal{M}u_0 = u_{F_{m, u_0}}$.

Analogously, there exist unique elements $u_{F_{g, u_0}}, u_{F_{k, u_0}} \in V$, so that $\mathcal{G}u_0 = u_{F_{g, u_0}}$ and $\mathcal{K}u_0 = u_{F_{k, u_0}}$, respectively, if we assume that $|g(u_0, v)| \leq c\|v\|_V$ and $|k(u_0, v)| \leq c\|v\|_V$ for all $v \in V$ and for each fixed $u_0 \in V$.

These boundedness conditions on m , g and k are satisfied thanks to the assumptions (IV), (V), (VI) and the embedding property (2). \square

Lemma 3.5. *The operators satisfy $\mathcal{M}^* = \mathcal{M}$, $\mathcal{G}^* = -\mathcal{G}$, $\mathcal{K}^* = \mathcal{K}$.*

Proof. The assertion follows from

$$\begin{aligned}\langle \mathcal{M}u, v \rangle_V &= m(u, v) = \overline{m(v, u)} = \overline{\langle \mathcal{M}v, u \rangle_V} = \overline{\langle v, \mathcal{M}^*u \rangle_V} = \langle \mathcal{M}^*u, v \rangle_V \\ \langle \mathcal{G}u, v \rangle_V &= g(u, v) = -\overline{g(v, u)} = -\overline{\langle \mathcal{G}v, u \rangle_V} = -\overline{\langle v, \mathcal{G}^*u \rangle_V} = \langle -\mathcal{G}^*u, v \rangle_V \\ \langle \mathcal{K}u, v \rangle_V &= k(u, v) = \overline{k(v, u)} = \overline{\langle \mathcal{K}v, u \rangle_V} = \overline{\langle v, \mathcal{K}^*u \rangle_V} = \langle \mathcal{K}^*u, v \rangle_V.\end{aligned}$$

□

From the definition of \mathcal{M}^* , \mathcal{G}^* , \mathcal{K}^* , we conclude that $\mathcal{M}^{**} = \mathcal{M}$, $\mathcal{G}^{**} = \mathcal{G}$, $\mathcal{K}^{**} = \mathcal{K}$, since $\langle \mathcal{M}u, v \rangle_V = \langle u, \mathcal{M}^*v \rangle_V = \overline{\langle \mathcal{M}^*v, u \rangle_V} = \overline{\langle v, \mathcal{M}^{**}u \rangle_V} = \langle \mathcal{M}^{**}u, v \rangle_V$ and, analogously, $\langle \mathcal{G}u, v \rangle_V = \langle \mathcal{G}^{**}u, v \rangle_V$ and $\langle \mathcal{K}u, v \rangle_V = \langle \mathcal{K}^{**}u, v \rangle_V$ for all $u, v \in V$.

We introduce the operator pencil $\mathcal{B}(\cdot) : \mathbb{C} \rightarrow \mathcal{L}(V, V)$ by

$$\mathcal{B}(\lambda) := \mathcal{K} - \lambda\mathcal{G} - \lambda^2\mathcal{M}, \quad \lambda \in \mathbb{C} \quad (3)$$

and define the adjoint operator pencil

$$\mathcal{B}^*(\lambda) := [\mathcal{B}(\lambda)]^* = \mathcal{K}^* - \bar{\lambda}\mathcal{G}^* - \bar{\lambda}^2\mathcal{M}^*, \quad \lambda \in \mathbb{C}.$$

Lemma 3.6 (Operator eigenvalue problem). *The quadratic eigenvalue problem (1) is equivalent to the operator eigenvalue problem for $\mathcal{B}(\cdot)$: Find $\lambda \in \mathbb{C}$, $u \in V \setminus \{0\}$, such that*

$$\mathcal{B}(\lambda)u = 0. \quad (4)$$

Proof. We know from the definition of \mathcal{M} , \mathcal{G} , \mathcal{K} that

$$\begin{aligned}\langle \mathcal{B}(\lambda)u, v \rangle_V &= \langle (\mathcal{K} - \lambda\mathcal{G} - \lambda^2\mathcal{M})u, v \rangle_V = \langle \mathcal{K}u, v \rangle_V - \lambda\langle \mathcal{G}u, v \rangle_V - \lambda^2\langle \mathcal{M}u, v \rangle_V \\ &= k(u, v) - \lambda g(u, v) - \lambda^2 m(u, v).\end{aligned}$$

Furthermore, $(\lambda, u) \in \mathbb{C} \times V$ is an eigenpair of problem (1) if and only if $k(u, v) - \lambda g(u, v) - \lambda^2 m(u, v) = 0$ for all $v \in V$, that is, $\langle \mathcal{B}(\lambda)u, v \rangle_V = 0$ for all $v \in V$. This is equivalent to $\mathcal{B}(\lambda)u = 0$, since $\langle \cdot, \cdot \rangle_V$ is positive definite, and therefore, (λ, u) is an eigenpair of problem (4). □

The assumptions (IV), (V), (VI) were made so that the operators \mathcal{M} and \mathcal{G} are compact (we will prove this in Section 3.3) and that the operator \mathcal{K} is invertible. It follows from Lemma 2.10 and Remark 2.6 that the operator $\mathcal{B}(\lambda)$ is a Fredholm operator with Index $\text{Ind } \mathcal{B}(\lambda) = 0$ for each $\lambda \in \mathbb{C}$.

Thanks to Remark 2.11, we know that $\mathcal{B}(\lambda)$ is invertible if and only if the equation $\mathcal{B}(\lambda)u = 0$ has only the trivial solution $u = 0$. Hence, we can write

$$\sigma(\mathcal{B}) = \{\lambda \in \mathbb{C} \mid \exists u \in V \setminus \{0\}, \text{ so that } \mathcal{B}(\lambda)u = 0\} = \{\lambda \in \mathbb{C} \mid \mathcal{B}(\lambda) \text{ is not invertible}\}.$$

3.3 Compactness of \mathcal{M} and \mathcal{G}

The aim of this subsection is the verification that the operators \mathcal{M} and \mathcal{G} are compact under the assumptions (I), (IV) and (V).

We consider any bounded sequence $\{u_n\}$ in V . Since V is compactly embedded into H due to assumption (I), there is a subsequence u_{n_k} which converges in H . For the sake of simplicity of notation, we denote this subsequence by $\{u_n\}$ and omit the subindex k . We conclude that $\{u_n\}$ is a Cauchy sequence in H . Assumption (IV) and the embedding property (2) imply that

$$\begin{aligned}\|\mathcal{M}u_n - \mathcal{M}u_m\|_V^2 &= |\langle \mathcal{M}u_n - \mathcal{M}u_m, \mathcal{M}u_n - \mathcal{M}u_m \rangle_V| = |m(u_n - u_m, \mathcal{M}u_n - \mathcal{M}u_m)| \\ &\leq c\|u_n - u_m\|_H \cdot \|\mathcal{M}u_n - \mathcal{M}u_m\|_H \\ &\leq c\|u_n - u_m\|_H \cdot \|\mathcal{M}u_n - \mathcal{M}u_m\|_V.\end{aligned}$$

Division by $\|\mathcal{M}u_n - \mathcal{M}u_m\|_V$ yields

$$\|\mathcal{M}u_n - \mathcal{M}u_m\|_V \leq c\|u_n - u_m\|_H.$$

Since $\{u_n\}$ is a Cauchy sequence in H , we obtain that $\{\mathcal{M}u_n\}$ is a Cauchy sequence in V . Exploiting that V is complete, we conclude that $\{\mathcal{M}u_n\}$ converges in V , which proves the compactness of \mathcal{M} .

By analogy with the operator \mathcal{M} , we introduce the operator $\mathcal{D} : V \rightarrow V$ by

$$\langle \mathcal{D}u, v \rangle_V = d(u, v) \quad \forall u, v \in V.$$

Then, with the same arguments as above, we get from assumption (V) that the operator \mathcal{D} is compact. Hence, the adjoint operator \mathcal{D}^* , which is given by

$$\langle \mathcal{D}u, v \rangle_V = \langle u, \mathcal{D}^*v \rangle_V \quad \forall u, v \in V,$$

is compact as well, see Werner (1997, Satz von Schauder (III.4.4)). Moreover, we have for all $u, v \in V$ that

$$\begin{aligned}\langle \mathcal{G}u, v \rangle_V &= g(u, v) = \overline{d(v, u)} - d(u, v) = \overline{\langle \mathcal{D}v, u \rangle_V} - \langle \mathcal{D}u, v \rangle_V \\ &= \overline{\langle v, \mathcal{D}^*u \rangle_V} - \langle \mathcal{D}u, v \rangle_V = \langle \mathcal{D}^*u, v \rangle_V - \langle \mathcal{D}u, v \rangle_V = \langle (\mathcal{D}^* - \mathcal{D})u, v \rangle_V.\end{aligned}$$

Therefore, by Remark 2.6, the operator $\mathcal{G} = \mathcal{D}^* - \mathcal{D}$ is compact.

3.4 Proof: $\lambda \in \sigma(\mathcal{B}) \implies -\bar{\lambda} \in \sigma(\mathcal{B})$

Returning to the eigenvalue problem for the operator pencil $\mathcal{B}(\cdot)$ and the Hamiltonian structure of the spectrum, we start with the proof that with $\lambda \in \mathbb{C}$ also $-\bar{\lambda} \in \mathbb{C}$ is an eigenvalue of problem (4).

Lemma 3.7. *The relation $\mathcal{B}^*(\lambda) = \mathcal{B}(-\bar{\lambda})$ is true for all $\lambda \in \mathbb{C}$.*

Proof. Each $\lambda \in \mathbb{C}$ satisfies $\mathcal{B}^*(\lambda) = \mathcal{K}^* - \bar{\lambda}\mathcal{G}^* - \bar{\lambda}^2\mathcal{M}^* = \mathcal{K} - (-\bar{\lambda})\mathcal{G} - (-\bar{\lambda})^2\mathcal{M} = \mathcal{B}(-\bar{\lambda})$. \square

Lemma 3.8. *The operators \mathcal{B} and \mathcal{B}^* have the same eigenvalues, that is, $\sigma(\mathcal{B}) = \sigma(\mathcal{B}^*)$.*

Proof. Since $\mathcal{B}(\lambda)$ is a Fredholm operator with index 0, we have that $\lambda \in \sigma(\mathcal{B})$ if and only if $\mathcal{B}(\lambda)$ is not invertible. By Remark 2.11, this is true if and only if $\mathcal{B}^*(\lambda)$ is not invertible, which is equivalent to $\lambda \in \sigma(\mathcal{B}^*)$. \square

Corollary 3.9. *If $\lambda \in \sigma(\mathcal{B})$ then $-\bar{\lambda} \in \sigma(\mathcal{B})$.*

It remains to show either that $\bar{\lambda} \in \sigma(\mathcal{B})$ or that $-\lambda \in \sigma(\mathcal{B})$ given $\lambda \in \sigma(\mathcal{B})$. The existence of the fourth eigenvalue that completes the Hamiltonian structure follows then immediately from Corollary 3.9.

3.5 Proof: $\lambda \in \sigma(\mathcal{B}) \implies \bar{\lambda} \in \sigma(\mathcal{B})$

The essential condition for this subsection is that the sesquilinear forms are real, see assumption (III). We use this property to evaluate the term $\mathcal{B}(\lambda)u$.

Lemma 3.10. *The relations $\overline{\mathcal{M}u} = \mathcal{M}\bar{u}$, $\overline{\mathcal{G}u} = \mathcal{G}\bar{u}$, $\overline{\mathcal{K}u} = \mathcal{K}\bar{u}$ are true for all $u \in V$.*

Proof. Assumption (III) implies for each $u \in V$ that

$$\langle \overline{\mathcal{M}u}, v \rangle_V = \overline{\langle \mathcal{M}u, \bar{v} \rangle_V} = \overline{m(u, \bar{v})} = m(\bar{u}, v) = \langle \mathcal{M}\bar{u}, v \rangle_V \quad \forall v \in V,$$

which proves the assertion for $\overline{\mathcal{M}u}$. The relations $\overline{\mathcal{G}u} = \mathcal{G}\bar{u}$ and $\overline{\mathcal{K}u} = \mathcal{K}\bar{u}$ follow analogously. \square

Corollary 3.11. *The relation $\overline{\mathcal{B}(\lambda)u} = \mathcal{B}(\bar{\lambda})\bar{u}$ is true for all $u \in V$.*

Proof. Lemma 3.10 yields for each $u \in V$ that

$$\overline{\mathcal{B}(\lambda)u} = \overline{(\mathcal{K} - \lambda\mathcal{G} - \lambda^2\mathcal{M})u} = \overline{\mathcal{K}u} - \overline{\lambda\mathcal{G}u} - \overline{\lambda^2\mathcal{M}u} = \mathcal{K}\bar{u} - \bar{\lambda}\mathcal{G}\bar{u} - \bar{\lambda}^2\mathcal{M}\bar{u} = \mathcal{B}(\bar{\lambda})\bar{u}.$$

\square

Corollary 3.12. *If $\lambda \in \mathbb{C}$ is an eigenvalue of $\mathcal{B}(\cdot)$ then so is $\bar{\lambda}$.*

Proof. If $\lambda \in \mathbb{C}$ is an eigenvalue of $\mathcal{B}(\cdot)$, then there is an element $u \in V \setminus \{0\}$, so that $\mathcal{B}(\lambda)u = 0$. Consequently, we have that $\overline{\mathcal{B}(\lambda)u} = 0$. Corollary 3.11 implies that $\mathcal{B}(\bar{\lambda})\bar{u} = 0$, that is, $\bar{\lambda}$ is an eigenvalue of $\mathcal{B}(\cdot)$, since $\bar{u} \in V \setminus \{0\}$. \square

This part of the proof holds even if $\mathcal{B}(\lambda)$ is not Fredholm, this means that the compact embedding part of assumption (I) and the special structure of g formulated in assumption (V) were not used here. Neither was condition (II) employed. The boundedness conditions in (IV)–(VI), however, can not be omitted, because we used them in the proof that the operators \mathcal{M} , \mathcal{G} and \mathcal{K} are well-defined, see Lemma 3.4.

3.6 The Hamiltonian eigenvalue symmetry

The following theorem is the consequence of the previous results.

Theorem 3.13 (Hamiltonian eigenvalue symmetry). *If the assumptions (I)–(VI) are satisfied, then, the spectrum of the quadratic eigenvalue problem (1),*

$$\lambda^2 m(u, v) + \lambda g(u, v) = k(u, v),$$

is symmetric with respect to the real and imaginary axes, that is, if $\lambda \in \mathbb{C}$ is an eigenvalue of problem (1), then so are $-\lambda$, $\bar{\lambda}$, $-\bar{\lambda}$.

The theorem is easily proven by the use of Corollary 3.9 ($\lambda \in \sigma(\mathcal{B}) \Rightarrow -\bar{\lambda} \in \sigma(\mathcal{B})$), Corollary 3.12 ($\lambda \in \sigma(\mathcal{B}) \Rightarrow \bar{\lambda} \in \sigma(\mathcal{B})$) and Lemma 3.6 which states that $\lambda \in \mathbb{C}$ is an eigenvalue of problem (1) if and only if $\mathcal{B}(\lambda)u = 0$.

Remark 3.14. *The same symmetry is given for non-linear eigenvalue problems of higher order,*

$$\sum_{i=0}^n \lambda^i m_i(u, v) = 0,$$

with alternating sequences of Hermitian and skew-Hermitian sesquilinear forms $m_i(\cdot, \cdot)$, if $-m_0$ satisfies condition (VI), the other Hermitian forms fulfil condition (IV), and (V) holds for the skew-Hermitian forms. Polynomial eigenvalue problems of this form were treated, for example, by Mehrmann and Watkins (2002).

4 Examples

Example 4.1. *Consider the counter-example of the introduction: $g(u, v) = \mathbf{i} m(u, v)$, $k(u, v) \equiv 0$. Obviously, k is not positive definite and the operator \mathcal{K} induced by $k(\cdot, \cdot)$ is the zero operator and therefore not invertible. Hence, the Fredholm theory cannot be applied.*

Note that, in general, we cannot conclude from this that the desired eigenvalue symmetry is not given; in this example, however, $\lambda = -\mathbf{i}$ is an eigenvalue, but $\lambda = \mathbf{i}$ is not.

Example 4.2. *Let V be a finite-dimensional complex Hilbert space; without loss of generality, $V = \mathbb{C}^n$, $n \in \mathbb{N}$. Let m, g, k be given by*

$$m(u, v) = \bar{v}^\top M u, \quad g(u, v) = \bar{v}^\top G u, \quad k(u, v) = \bar{v}^\top K u \quad \forall u, v \in V,$$

where M, G, K are complex $n \times n$ -matrices.

Let $\langle \cdot, \cdot \rangle_V$ denote the usual inner product, $\langle u, v \rangle_V := \bar{v}^\top u$. The operator \mathcal{M} induced by $m(u, v) = \langle \mathcal{M}u, v \rangle_V$ for all $u, v \in V$ is then defined by $m(u, v) = \bar{v}^\top \mathcal{M}u$, that is, $\mathcal{M} = M$. Analogously, we have that $\mathcal{G} = G$ and $\mathcal{K} = K$. Hence, the eigenvalue problem is equivalent to the matrix eigenvalue problem: Find $\lambda \in \mathbb{C}$, $u \in \mathbb{C}^n$, such that

$$\lambda^2 M u + \lambda G u = K u.$$

The assumptions (II) and (III) imply that M , G and K are real matrices with $M = M^\top$, $G = -G^\top$ and $K = K^\top$.

Indeed, the Hamiltonian eigenvalue symmetry can be proven then: Assuming that λ is an eigenvalue, we get by conjugating and transposing the equation that $-\bar{\lambda}$ is an eigenvalue, too. The characteristic polynomial $\det(\lambda^2 M + \lambda G - K)$ has real coefficients, which implies that $\bar{\lambda}$ is an eigenvalue as well, and therefore so is $-\lambda$.

Note that (VI), demanding the invertibility of K , needs not to be satisfied, here; nor are (IV) or (V) essential. Nevertheless, it is necessary that M , G and K are real. Indeed, one easily finds complex counter-examples, compare $M = I$, $G = 2iI$, $K = 2I$, where $\lambda = i + 1$ and $-\bar{\lambda} = i - 1$ are eigenvalues, but $\bar{\lambda}$ and $-\lambda$ are not.

Finally, let us consider a quadratic eigenvalue problem that arises in numerical practice. In the neighbourhood of polyhedral corners, the singular part of the solution to the linear elasticity problem has a special structure. The computation of the corner singularities can be reduced to a quadratic eigenvalue problem of type (1), which we learnt from Leguillon (1995). Since the underlying problem is an elliptic boundary value problem, Kozlov, Maz'ya, and Roßmann (2000) suggested to apply a Mellin transformation from which the Hamiltonian eigenvalue structure can be obtained as well. Further details including the operator eigenvalue problem formulated in Section 3.2 are given by Apel, Sändig, and Solov'ev (2002b).

Example 4.3. *The study of corner singularities leads to the consideration of a ball circumscribing the polyhedral corner. Using spherical coordinates and a specific approach for the structure of the solution to the linear elasticity, we obtain the eigenvalue problem*

$$\lambda^2 m(u, v) + \lambda g(u, v) = k(u, v)$$

with $\lambda \in \mathbb{C}$, $u, v \in V$, where $V = H_0^1(\Omega)^3$ and $H = L^2(\Omega)^3$ are spaces of vector functions over a subdomain Ω of the unit sphere in \mathbb{R}^3 . This eigenvalue problem was written down in a slightly different form by Leguillon (1995); we refer also to the work by Apel, Sändig, and Solov'ev (2002b), which was the basis for the derivation of the Hamiltonian eigenvalue symmetry in this paper.

The space $H_0^1(\Omega)$ is compactly embedded into $L^2(\Omega)$, hence so is V into H . The assumptions on m , g , k are satisfied in the given problem; therefore the suggested symmetry of the spectrum can be concluded, as it was done rather shortly by Apel, Sändig, and Solov'ev (2002b, Theorem 10) or Leguillon (1995) or only for the discrete formulation, for instance, by Apel, Mehrmann, and Watkins (2002a, Proposition 1), see also Mehrmann and Watkins (2002) and references therein.

5 Summary

We studied a quadratic eigenvalue problem and formulated conditions under which the Hamiltonian eigenvalue symmetry can be proven. To derive the Hamiltonian structure, we

transformed the problem into an eigenvalue problem for an operator pencil and applied Fredholm theory. Since it is usually difficult to check whether a given operator is a Fredholm operator with index 0 or not, we formulated simplified conditions in (I)–(VI) which imply the Fredholm property. At last, we proved that under these assumptions the spectrum of the quadratic eigenvalue problem

$$\lambda^2 m(u, v) + \lambda g(u, v) = k(u, v)$$

has the desired structure. In general, these conditions are easy to check, once the functions m , g , k are given.

Finally, we presented an application and discussed the necessity of the assumptions in two further examples.

Acknowledgement

For fruitful discussion and valuable advice, special thanks go to Thomas Apel (Universität der Bundeswehr München), Bernd Silbermann (Technische Universität Chemnitz) and Sergey Solov'ev (Kazan State University). We also thank the DFG (Deutsche Forschungsgemeinschaft) for financial support.

References

- Alt, H. W., 1999. *Lineare Funktionalanalysis*. Springer, Berlin, Heidelberg, New York, 3. Auflage.
- Apel, T., Mehrmann, V., Watkins, D., 2002a. Structured eigenvalue methods for the computation of corner singularities in 3D anisotropic elastic structures. *Comput. Methods Appl. Mech. Engrg.* 191, 4459–4473.
- Apel, T., Sändig, A.-M., Solov'ev, S. I., 2002b. Computation of 3d vertex singularities for linear elasticity: Error estimates for a finite element method on graded meshes. *Math. Model. Numer. Anal.* 36, 1043–1070.
- Bai, Z., 2002. Krylov subspace techniques for reduced-order modeling of large-scale dynamical systems. *Appl. Numer. Math.* 43, 9–44.
- Benner, P., Faßbender, H., 1997. An implicitly restarted symplectic Lanczos method for the Hamiltonian eigenvalue problem. *Lin. Alg. Appl.* 263, 75–111.
- Benner, P., Faßbender, H., 2000. An implicitly restarted symplectic Lanczos method for the symplectic eigenvalue problem. *SIAM Journal on Matrix Analysis and Applications* 22 (3), 682–713.
- Cotlar, M., Cignoli, R., 1974. *An Introduction to Functional Analysis*. North-Holland Publishing Company, Amsterdam.

- Freund, R. W., 1994. Lanczos-type algorithms for structured non-Hermitian eigenvalue problems. SIAM, pp. 243–245, proceedings of the Cornelius Lanczos International Centenary Conference.
- Kozlov, V. A., Maz'ya, V. G., Roßmann, J., 2000. Spectral problems associated with corner singularities of solutions to elliptic equations. *Mathematical Surveys and Monographs*, Volume 85, American Mathematical Society.
- Leguillon, D., 1995. Computation of 3d singularities in elasticity. In: Costabel, M., Dauge, M., Nicaise, S. (Eds.), *Boundary value problems and integral equations in nonsmooth domains*. Vol. 167 of *Lecture notes in pure and applied mathematics*. Marcel Dekker, New York, pp. 161–170, proceedings of a conference at CIRM, Luminy, France, May 3-7, 1993.
- Mehrmann, V., Watkins, D., 2001. Structure-preserving methods for computing eigenpairs of large sparse skew-Hamiltonian/Hamiltonian pencils. *SIAM J. on Sci. Comp.* 22 (6), 1905–1925.
- Mehrmann, V., Watkins, D., 2002. Polynomial eigenvalue problems with Hamiltonian structure. *Electron. Trans. Numer. Anal.* 13, 106–118.
- Prößdorf, S., 1974. *Einige Klassen singulärer Gleichungen*. Akademie-Verlag, Berlin.
- Schiehlen, W., 1990. *Multibody Systems Handbook*. Springer-Verlag.
- Watkins, D., 2002. *Fundamentals of Matrix Computations*. Wiley.
- Watkins, D., 2004. On Hamiltonian and symplectic Lanczos processes. *Lin. Alg. Appl.* 385, 23–45.
- Werner, D., 1997. *Funktionalanalysis*. Springer, Berlin, Heidelberg, New York, 2. Auflage.