

Nonlinear Optimization Exercises 6

1. Show for the Line-Search Newton-CG method that the generated step directions are descent directions.
2. Prove: In the Trust-Region Steihaug-CG method the length of the step is nondecreasing. In the notation of the algorithm, $0 = \|p_0\| < \dots < \|p_j\| < \dots < \|p\| = \Delta$.
3. Nonlinear Systems of Equations: Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be of the form

$$F(x) = \begin{pmatrix} x^T A_1 x + b_1^T x + c_1 \\ x^T A_2 x + b_2^T x + c_2 \end{pmatrix}.$$

with $A_i \in \mathbb{R}^{2 \times 2}$ symmetric, $b_i \in \mathbb{R}^2$, $c_i \in \mathbb{R}$ for $i \in \{1, 2\}$ and Jacobian $J(x)$. Denote the set of degenerate (non regular) points by $D = \{x \in \mathbb{R}^2 : J(x) \text{ singular}\}$. If possible, give a choice of coefficients A_i , b_i , c_i so that

- (a) $D = \emptyset$
- (b) $D = \mathbb{R}^2$
- (c) $|D| = 1$
- (d) $D = \{x : a_1^T x = 0\} \cup \{x : a_2^T x = 0\}$ for given $a_1, a_2 \in \mathbb{R}^2$.

4. Consider the optimization problem

$$\begin{array}{ll} \min & c^T x \\ \text{s.t.} & \frac{1}{2} x^T A x \leq b_1 \\ & a^T x \leq b_2 \\ & x \in \mathbb{R}^3 \end{array} \quad \text{where } c = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, a = \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}, b = \begin{pmatrix} \frac{1}{2} \\ -1 \end{pmatrix}.$$

Give the set of optimal solutions. Describe the set of limiting directions for the point $\bar{x} = (1, 0, 0)$, the linearised tangent cone $F_1(\bar{x})$, and its consequences on the existence of Lagrange multipliers. Provide a description of the feasible set, so that Lagrange-multipliers exist for all optimal solutions.

5. A mathematical program with equilibrium constraints (MPEC) is an optimization problem of the form

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & g(x) \leq 0, h(x) = 0, \\ & G(x) \geq 0, H(x) \geq 0, G(x)^T H(x) = 0 \end{array} \quad \begin{array}{l} f : \mathbb{R}^n \rightarrow \mathbb{R} \\ \text{with } g : \mathbb{R}^n \rightarrow \mathbb{R}^m, h : \mathbb{R}^n \rightarrow \mathbb{R}^p \\ G, H : \mathbb{R}^n \rightarrow \mathbb{R}^M \end{array}$$

Prove that for an MPEC there is no feasible point in which LICQ holds.