

# Stability and efficiency of a recursive multirate Rosenbrock method

Karen Kuhn<sup>1</sup>   Jens Lang<sup>2</sup>

Many physical phenomena contain different time scales. One way to solve the descriptive PDE is to discretize first in space and then apply a normal singlerate time integrator to the resulting ODE system. For problems with different time scales this might end up in very small time steps which have to be applied also to components with much less activity. That is why the application of multirate methods is reasonable. Different time step sizes are used for different components, depending on their individual activity. Since the stability character of a singlerate method usually is not carried over to the corresponding multirate method, we study the asymptotic stability for several multirate Rosenbrock methods. For several test problems we compare the needed CPU time of the multirate and the respective singlerate methods.

**Acknowledgment:** This work is supported by the “Excellence Initiative” of the German Federal and State Governments and the Graduate School of Computational Engineering at Technische Universität Darmstadt.

References:

- [1] Savcenco, V. and Hundsdorfer, W.H. and Verwer, J.G., *A multirate time stepping strategy for stiff ODEs*, BIT, 47, 2007.
- [2] Kuhn, K. and Lang, J., *Comparison of asymptotic stability for recursive ROS-multirate methods*, TU Darmstadt, Preprint 2625, 2011, submitted to the Journal of Comp. and Appl. Math. 2012.

---

<sup>1</sup> Technische Universität Darmstadt, Graduate School CE, Dolivostraße 15, D-64293 Darmstadt, [kuhn@gsc.tu-darmstadt.de](mailto:kuhn@gsc.tu-darmstadt.de)

<sup>2</sup> Technische Universität Darmstadt, [lang@mathematik.tu-darmstadt.de](mailto:lang@mathematik.tu-darmstadt.de)