

Odd characteristic classes in entire cyclic homology and equivariant loop space homology

Sergio Cacciatori¹ and Batu Güneysu²

¹ Dipartimento di Scienza e Alta Tecnologia, Università dell'Insubria, via Valleggio 11, 22100, Como, Italy and INFN - Sezione di Milano, via Celoria 16, 20133, Milano, Italy
email: sergio.cacciatori@uninsubria.it

² Mathematisches Institut, Universität Bonn, Endenicher Allee 60, 53115 Bonn, Germany
email: gueneysu@math.uni-bonn.de

Abstract

Given a compact manifold M and a smooth map $g : M \rightarrow U(l \times l; \mathbb{C})$ from M to the Lie group of unitary $l \times l$ matrices with entries in \mathbb{C} , we construct a Chern character $\text{Ch}^-(g)$ which lives in the odd part of the equivariant (entire) cyclic Chen-normalized cyclic complex $\mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$ of M , and which is mapped to the odd Bismut-Chern character under the equivariant Chen integral map. It is also shown that the assignment $g \mapsto \text{Ch}^-(g)$ induces a well-defined group homomorphism from the K^{-1} theory of M to the odd homology group of $\mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$.

Let M be a closed Riemannian spin manifold with its Clifford multiplication

$$c : \Omega(M) \longrightarrow \text{End}(S)$$

and its Dirac operator D acting in $L^2(M, S)$, and given $g \in C^\infty(M, U(l \times l; \mathbb{C}))$ let D_g denote the twisted Dirac operator

$$D_g := g^{-1}Dg = D + c(g^{-1}dg),$$

considered to be acting on $L^2(M, S \otimes \mathbb{C}^l)$. Then with

$$D_{g,s} := (1-s)D + sD_g, \quad s \in [0, 1],$$

the odd dimensional variant of Atiyah-Singer's 'index' theorem states that if M is odd dimensional, then [8]

$$\frac{1}{2\pi} \int_0^1 \text{Tr} \left[\dot{D}_{g,s} \exp(-D_{g,s}^2) \right] ds = \int_M \hat{A}(M) \wedge \text{ch}^-(g), \quad (1)$$

where $\text{ch}^-(g) \in \Omega^-(M)$ denotes the odd Chern character. The left hand side of (1) is precisely the spectral flow $\text{sf}(D, D_g)$ [8]. Furthermore, on the RHS of this formula, the odd Chern character can be obtained integration along the fiber of $M \times I \rightarrow M$ of the even Chern character of an appropriately chosen connection on $M \times I$ [8]. In fact, this formula can be proved by noting the LHS admits an infinite dimensional version of such an even/odd periodicity [4, 5] in terms of the eta form.

Being motivated by the considerations of Atiyah and Bismut [1, 2] for the even-dimensional case, one finds that another very elegant and geometric, however purely formal, way to prove (1) is to assume the existence of a Duistermaat-Heckmann localization formula for the smooth loop space LM : indeed, the spin structure on M induces an orientation on LM [1] and the path integral formalism entails the elegant, however mathematically ill-defined, formula (the even-dimensional variant of this formula is well-known [2] and the odd-dimensional case can be proved similarly [14])

$$\frac{1}{2\pi} \int_0^1 \text{Tr} \left[\dot{D}_{g,s} \exp(-D_{g,s}^2) \right] ds = \int_{LM} \exp(-\beta) \wedge \text{Bch}^-(g), \quad (2)$$

where $\beta = \beta_0 + \beta_2 \in \Omega^+(LM)$ denotes the even differential form defined on smooth vector fields X, Y on LM by

$$\beta_0(X) := \int_0^1 |X_s|^2 ds, \quad \beta_2(X, Y) := \int_0^1 (\nabla X_s / \nabla s, Y_s) ds,$$

and where $\text{Bch}^-(g) \in \Omega^-(M)$ denotes the odd Bismut-Chern character [3, 18]. Now both differential forms $\exp(-\beta)$ and $\text{Bch}^-(g)$ are equivariantly closed (cf. Section 4 for the definition of the degree -1 differential P),

$$(d + P) \exp(-\beta) = 0 = (d + P) \text{Bch}^-(g)$$

and so is their product. As the fixed point set of the \mathbb{T} -action on LM given by rotating every loop is precisely $M \subset LM$, a hypothetical Duistermaat-Heckmann localization formula immediately gives

$$\int_{LM} \exp(-\beta) \wedge \text{Bch}^-(g) = \int_M \hat{A}(M) \wedge \exp(-\beta)|_M \wedge \text{Bch}^-(g)|_M,$$

as $\hat{A}(M)$ is the inverse of the (appropriately renormalized) Euler class of the normal bundle of $M \subset LM$. This proves (1), as clearly $\exp(-\beta)|_M = 1$ and by construction $\text{Bch}^-(g)|_M = \text{ch}^-(g)$.

A direct implementation of the above arguments is not possible, as the right hand side of formula (2) is not well-defined for various reasons. For example, there exists no volume measure on LM , while smooth loops have Wiener measure zero, and, on the other hand, it is notoriously difficult to produce a variant of the super complex $(\Omega(LM), d + P)$ if one replaces LM with the smooth Banach manifold of *continuous loops*. Nevertheless and strikingly, the above formal manipulations lead to the powerful machinery of hypoelliptic Dirac and Laplace operators, as is explained in [3] and the references therein.

However, a possible way out of these problems has been proposed by Getzler, Jones and Petrack (GJP) [11] [9]. In this approach, the idea is to take as model for $\Omega(LM)$ the space of equivariant Chen integrals: these are given by the image of a morphism of super complexes (cf. Section 4 below for the relevant definitions)

$$\rho : (\mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T})), b + B) \longrightarrow (\widehat{\Omega}(LM), d + P).$$

Above, $\mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$ denotes the Chen-normalized entire cyclic (or Connes) complex of the locally convex unital DGA $\Omega_{\mathbb{T}}(M \times \mathbb{T})$, and $\widehat{\Omega}(LM)$ denotes a completed space of smooth differential forms on LM . Now the GJP-program for infinite dimensional localization is as follows: here it is conjectured that the composition

$$\int_{LM} \exp(-\beta) \wedge \rho(\cdot) : \mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T})) \longrightarrow \mathbb{C},$$

is a mathematically well-defined continuous functional, and that

- $\int_{LM} \exp(-\beta) \wedge \rho(\cdot)$ is odd (as LM is formally odd-dimensional if M is so [3]) and co-closed, meaning that it vanishes on the exact elements of $\mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$,
- if $w \in \mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$ is closed, then one has the 'Duistermaat-Heckmann localization formula'

$$\int_{LM} \exp(-\beta) \wedge \rho(w) = \int_M \hat{A}(TM) \wedge \rho(w)|_M.$$

If in addition one could canonically construct an element

$$\text{Ch}^-(g) \in \mathcal{N}_\epsilon^-(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$$

such that

- i) $\text{Ch}^-(g)$ is closed
- ii) $\rho(\text{Ch}^-(g)) = \text{Bch}^-(g)$
- iii) $\rho(\text{Ch}^-(g))|_M = \text{ch}^-(g)$,

then from the above observations we would immediately obtain a proof of (1) within the GJP-program for infinite dimensional localization. Note that in the even dimensional case such a Chern character has been constructed as an even cycle in $\mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$ in [11].

The aim of this paper is precisely to construct a canonically given element

$$\text{Ch}^-(g) \in \mathcal{N}_\epsilon^-(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$$

satisfying the above properties i), ii), iii). In fact, our main results Theorem 5.1 and Theorem 5.4 below construct $\text{Ch}^-(g)$ for M a compact manifold (possibly with boundary), which satisfies i) and iii) and in addition ii) if M is closed (so that LM is a well-defined smooth Fréchet manifold). We also show in Theorem 5.1 that the assignment $g \mapsto \text{Ch}^-(g)$ induces a well-defined group homomorphism

$$\mathbb{K}^{-1}(M) \longrightarrow \mathcal{N}(\Omega_{\mathbb{T}}(M \times \mathbb{T})).$$

Finally, taking for granted that the even variant of $\text{Ch}^-(g)$ and $\text{BCh}^-(g)$ have been previously defined [11, 2], we establish an even/odd periodicity, relating these constructions to ours, showing another analogy to (1).

Acknowledgements: The authors would like to thank Jean-Michel Bismut, Markus Pflaum and Shu Shen for their discussions. We are very grateful to Matthias Ludewig for sharing his construction of the equivariant Chen integral map with us.

1 Cyclic bar complex of a differential graded algebra (DGA)

In the sequel, we understand all our linear spaces to be over \mathbb{C} . Assume we are given a unital DGA Ω , that is,

- Ω is a unital algebra
- $\Omega = \bigoplus_{j=-\infty}^{\infty} \Omega^j$ is graded into subspaces $\Omega^j \subset \Omega$ such that $\Omega^i \Omega^j \subset \Omega^{i+j}$ for all $i, j \in \mathbb{Z}$, there is a degree +1 differential $d : \Omega \rightarrow \Omega$ which satisfies the graded Leibnitz rule.

Note that the space $\underline{\Omega} := \Omega / (\mathbb{C} \cdot \mathbf{1})$ is a graded linear space (but not canonically an algebra), and the space of cyclic chains $\mathcal{C}(\Omega)$ is defined as

$$\mathcal{C}(\Omega) := \bigoplus_{n=0}^{\infty} \Omega \otimes \underline{\Omega}^{\otimes n}.$$

We give $\Omega \otimes \underline{\Omega}^{\otimes n}$ the grading

$$\Omega \otimes \underline{\Omega}^{\otimes n} = \bigoplus_{j=0}^{\infty} \bigoplus_{j_0+\dots+j_n=j-n} \Omega^{j_0} \otimes \underline{\Omega}^{j_1} \otimes \dots \otimes \underline{\Omega}^{j_n},$$

which induces a linear map

$$\Gamma : \mathcal{C}(\Omega) \longrightarrow \mathcal{C}(\Omega), \quad \Gamma(w_0, w_1, \dots) := ((-1)^{\deg(w_0)}w_0, (-1)^{\deg(w_1)}w_1, \dots).$$

Since we have $\Gamma^2 = 1$, we can define a superstructure $\mathcal{C}(\Omega) = \mathcal{C}^+(\Omega) \oplus \mathcal{C}^-(\Omega)$ by setting

$$\mathcal{C}^{\pm}(\Omega) := \{w \in \mathcal{C}(\Omega) : \Gamma w = \pm w\}.$$

The following notation will be useful in the sequel:

Notation 1.1. Given $a \in \Omega \otimes \underline{\Omega}^{\otimes n}$ we define

$$\langle a \rangle := (\dots, a, \dots) \in \mathcal{C}(\Omega)$$

to be the cochain which has a in its n -th slot and 0 anywhere else.

We have the Hochschild map of the DGA-category

$$b : \mathcal{C}(\Omega) \longrightarrow \mathcal{C}(\Omega)$$

defined on $\Omega^{j_0} \otimes \underline{\Omega}^{j_1} \otimes \dots \otimes \underline{\Omega}^{j_n}$ by

$$\begin{aligned} b \langle \omega_0 \otimes \dots \otimes \omega_n \rangle &= \langle d\omega_0 \otimes \dots \otimes \omega_i \otimes \dots \otimes \omega_n \rangle \\ &\quad - \sum_{i=1}^n (-1)^{j_0+\dots+j_{i-1}-i+1} \langle \omega_0 \otimes \dots \otimes d\omega_i \otimes \dots \otimes \omega_n \rangle \\ &\quad - \sum_{i=0}^{n-1} (-1)^{j_0+\dots+j_i-i} \langle \omega_0 \otimes \dots \otimes \omega_i \omega_{i+1} \otimes \dots \otimes \omega_n \rangle \\ &\quad + (-1)^{(j_n-1)(j_0+\dots+j_{n-1}-n+1)} \langle (\omega_n \omega_0) \otimes \omega_1 \otimes \dots \otimes \omega_{n-1} \rangle, \end{aligned}$$

and Connes' operator

$$B : \mathcal{C}(\Omega) \longrightarrow \mathcal{C}(\Omega),$$

which is defined on $\Omega^{j_0} \otimes \underline{\Omega}^{j_1} \otimes \dots \otimes \underline{\Omega}^{j_n}$ by

$$B \langle \omega_0 \otimes \dots \otimes \omega_n \rangle = \sum_{i=0}^n (-1)^{(r_{i-1}+1)(r_n-r_{i-1})} \langle 1 \otimes \omega_i \otimes \dots \otimes \omega_n \otimes \omega_0 \otimes \dots \otimes \omega_{i-1} \rangle,$$

with $r_l = j_0 + \dots + j_l - l$. It is a well-known fact that one has

$$b^2 = 0, \quad B^2 = 0, \quad bB + Bb = 0, \quad \Gamma b = -\Gamma b, \quad \Gamma B = -\Gamma B.$$

We get the super complex

$$\mathcal{C}^+(\Omega) \xrightarrow{b+B} \mathcal{C}^-(\Omega) \xrightarrow{b+B} \mathcal{C}^+(\Omega). \quad (3)$$

The subspace $\mathcal{D}(\Omega) \subset \mathcal{C}(\Omega)$ is defined to be the linear span of all $w \in \mathcal{C}(\Omega)$ that satisfy one of the following relations:

- for all $n \in \mathbb{N}$ there exists $1 \leq r \leq n$, $f \in \Omega^0$, $\omega_0 \in \Omega$, $\omega_s \in \underline{\Omega}$, $s \neq r$, with

$$\langle w_n \rangle = \langle \omega_0 \otimes \cdots \otimes \omega_{r-1} \otimes f \otimes \omega_{r+1} \otimes \cdots \otimes \omega_n \rangle. \quad (4)$$

- for all $n \in \mathbb{N}$ there exists $1 \leq r \leq n$, $f \in \Omega^0$, $\omega_0 \in \Omega$, $\omega_s \in \underline{\Omega}$, $s \neq r$, with

$$\begin{aligned} & \langle \omega_0 \otimes \cdots \otimes \omega_{r-1} f \otimes \omega_{r+1} \otimes \cdots \otimes \omega_n \rangle + \langle \omega_0 \otimes \cdots \otimes \omega_{r-1} \otimes df \otimes \omega_{r+1} \otimes \cdots \otimes \omega_n \rangle \\ & - \langle \omega_0 \otimes \cdots \otimes \omega_{r-1} \otimes f \omega_{r+1} \otimes \cdots \otimes \omega_n \rangle. \end{aligned} \quad (5)$$

The maps Γ, b, B map $\mathcal{D}(\Omega)$ to itself, so that with

$$\mathcal{D}^\pm(\Omega) := \{w \in \mathcal{D}(\Omega) : \Gamma w = \pm w\},$$

there is a super complex

$$\mathcal{D}^+(\Omega) \xrightarrow{b+B} \mathcal{D}^-(\Omega) \xrightarrow{b+B} \mathcal{D}^+(\Omega).$$

With $\mathcal{N}^\pm(\Omega) := \mathcal{C}^\pm(\Omega)/\mathcal{D}^\pm(\Omega)$, the induced quotient complex

$$\mathcal{N}^+(\Omega) \xrightarrow{b+B} \mathcal{N}^-(\Omega) \xrightarrow{b+B} \mathcal{N}^+(\Omega).$$

Whenever there is no danger of confusion, the equivalence class of $w \in \mathcal{C}(\Omega)$ in $\mathcal{N}(\Omega)$ is denoted with the same symbol again.

2 Entire cyclic homology of a locally convex unital DGA

We recall that a topological vector space is called locally convex, if the topology is induced by a family of seminorms, noting that then the topology is equivalent to the topology induced by all continuous seminorms.

Definition 2.1. By a locally convex unital DGA we understand a unital DGA Ω which is also a locally convex Hausdorff space, such that

- the differential is continuous, e.g., for every continuous seminorm ε on Ω there exists a continuous seminorm ε' on Ω such that

$$\varepsilon(dw) \leq \varepsilon'(\omega) \quad \text{for all } \omega \in \Omega \quad (6)$$

- the multiplication is jointly continuous, e.g., for every continuous seminorm ε on Ω there exists a continuous seminorm ε' on Ω such that

$$\varepsilon(\omega_1 \omega_2) \leq \varepsilon'(\omega_1) \varepsilon'(\omega_2) \quad \text{for all } \omega_1, \omega_2 \in \Omega. \quad (7)$$

The space $\underline{\Omega}$ becomes a graded locally convex Hausdorff space, and we equip the algebraic tensor product $\underline{\Omega} \otimes \underline{\Omega}^{\otimes n}$ with the induced family of π -tensor seminorms, that is,

$$\varepsilon_n(\omega) = \inf \left\{ \sum_{\alpha} \varepsilon(\omega_0^{(1)}) \cdots \varepsilon(\omega_n^{(\alpha)}) : \omega = \sum_{\alpha} \omega_0^{(\alpha)} \otimes \cdots \otimes \omega_n^{(\alpha)} \right\},$$

where the sum runs through all representations of ω as a finite sum of elementary tensors, and where ε is a continuous seminorm on $\underline{\Omega}$.

Definition 2.2. The space of *entire cyclic chains* $\mathcal{C}_\varepsilon(\Omega)$ is defined to be the closure of $\mathcal{C}(\Omega)$ with respect to the seminorms

$$\kappa_\varepsilon(w) := \sum_{n=0}^{\infty} \frac{\varepsilon_n(w_n)}{\sqrt{n!}},$$

where ε is an arbitrary continuous seminorm on Ω .

The space $\mathcal{C}_\varepsilon(\Omega)$ is a complete locally convex Hausdorff space. Note that the above family of seminorms is equivalent to the family of seminorms

$$\kappa_{\varepsilon,l}(w) := \sum_{n=0}^{\infty} \frac{\varepsilon_n(w_n) l^n}{\sqrt{n!}} < \infty,$$

where ε is an arbitrary continuous seminorm on Ω and $l \in \mathbb{N}$, as $l\varepsilon$ is again a continuous seminorm and the ε_n 's are cross semi-norms. Thus, our growth conditions are modelled on the entire growth conditions for ungraded Banach algebras by Getzler/Szenes from [12].

Before stating the next auxiliary result, we recall that a continuous linear map from a locally convex Hausdorff space \mathcal{X} to a complete locally convex Hausdorff space \mathcal{Y} can be uniquely extended to a continuous linear map $\hat{\mathcal{X}} \rightarrow \mathcal{Y}$, noting that the completion $\hat{\mathcal{X}}$ is Hausdorff again. This can be proved precisely as for normed spaces.

Lemma 2.3. *The operators Γ, b, B map $\mathcal{C}(\Omega)$ continuously to itself, in particular, with*

$$\mathcal{C}_\varepsilon^\pm(\Omega) := \{w \in \mathcal{C}_\varepsilon(\Omega) : \Gamma w = \pm w\},$$

there is a well-defined super complex

$$\mathcal{C}_\varepsilon^+(\Omega) \xrightarrow{b+B} \mathcal{C}_\varepsilon^-(\Omega) \xrightarrow{b+B} \mathcal{C}_\varepsilon^+(\Omega). \quad (8)$$

Proof. Let ε be an arbitrary continuous seminorm on Ω . Clearly, one has $\kappa_\varepsilon(\Gamma w) \leq \kappa_\varepsilon(w)$ for all $w \in \mathcal{C}(\Omega)$.

Pick continuous seminorms $\varepsilon', \varepsilon''$ on Ω such that for all $\omega \in \Omega$ one has $\varepsilon(d\omega) \leq \varepsilon''(\omega)$ and such that for all $\omega_1, \omega_2 \in \Omega$ one has $\varepsilon(\omega_1 \omega_2) \leq \varepsilon'(\omega_1) \varepsilon'(\omega_2)$. Using $n+1 \leq 2^n$ it is then easily checked that

$$\kappa_\varepsilon(bw) \leq C \max(\kappa_{\varepsilon'}, \kappa_{\varepsilon''})(w) \quad \text{for all } w \in \mathcal{C}(\Omega).$$

Likewise, it follows immediately that $\kappa_\varepsilon(Bw) \leq C \kappa_\varepsilon(w)$ for all $w \in \mathcal{C}(\Omega)$. ■

Defining the subspace $\mathcal{D}_\epsilon(\Omega) \subset \mathcal{C}_\epsilon(\Omega)$ as the closure of $\mathcal{D}(\Omega)$, it follows automatically that the maps Γ, b, B map $\mathcal{D}(\Omega)$ continuously to itself, too, producing with

$$\mathcal{N}_\epsilon^\pm(\Omega) := \mathcal{C}_\epsilon^\pm(\Omega) / \mathcal{D}_\epsilon^\pm(\Omega)$$

the quotient complex

$$\mathcal{N}_\epsilon^+(\Omega) \xrightarrow{b+B} \mathcal{N}_\epsilon^-(\Omega) \xrightarrow{b+B} \mathcal{N}_\epsilon^+(\Omega). \quad (9)$$

Finally we can give:

Definition 2.4. The complex (8) is called the (reduced) *entire cyclic complex* of Ω and its homology groups are denoted with $\mathrm{HC}_\epsilon^\pm(\Omega)$. Likewise, the complex (9) is called the (reduced) *Chen-normalized entire cyclic complex* of Ω and its homology groups are denoted with $\mathrm{HN}_\epsilon^\pm(\Omega)$.

Above, 'reduced' refers to the fact that we work with $\Omega \otimes \underline{\Omega}^{\otimes n}$ rather than $\Omega^{\otimes(n+1)}$, which leads to a simpler formula for the Connes differential B .

3 The unital locally convex DGA $\Omega_{\mathbb{T}}(N \times \mathbb{T})$

Assume N is a manifold (possibly with boundary) and denote with \mathbb{T} the 1-sphere. We denote by $\Omega_{\mathbb{T}}(N \times \mathbb{T})$ the smooth \mathbb{T} -invariant differential forms on $N \times \mathbb{T}$, where \mathbb{T} acts trivially on N and by rotation on itself. Every element of $\Omega_{\mathbb{T}}(N \times \mathbb{T})$ can be uniquely written in the form $\alpha + \vartheta_{\mathbb{T}} \wedge \beta$ for some $\alpha, \beta \in \Omega(N)$, where $\vartheta_{\mathbb{T}}$ denotes the canonical 1-form on \mathbb{T} . We turn $\Omega_{\mathbb{T}}(N \times \mathbb{T})$ into a unital algebra by means of $\Omega_{\mathbb{T}}(N \times \mathbb{T}) \subset \Omega(N \times \mathbb{T})$, and give $\Omega_{\mathbb{T}}(N \times \mathbb{T})$ the grading

$$\alpha + \vartheta_{\mathbb{T}} \wedge \beta \in \Omega_{\mathbb{T}}^j(N \times \mathbb{T}) \iff \alpha \in \Omega^j(N), \beta \in \Omega^{j+1}(N).$$

With $\partial_{\mathbb{T}}$ the canonical vector field on \mathbb{T} , we have the differential $d_{\mathbb{T}} = d + \iota_{\partial_{\mathbb{T}}}$ defined by

$$d_{\mathbb{T}}(\alpha + \vartheta_{\mathbb{T}} \wedge \beta) = d\alpha + \beta - \vartheta_{\mathbb{T}} \wedge d\beta, \quad \text{if } \alpha + \vartheta_{\mathbb{T}} \wedge \beta \text{ is homogeneous,}$$

finally turning $\Omega_{\mathbb{T}}(N \times \mathbb{T})$ into a unital DGA.

Remark 3.1. Given a manifold X (possibly with boundary), the wedge product and the de Rham differential is continuous with respect to the canonical locally convex structure on $\Omega(X)$ [17]. In addition, if B is a vector field on X then the contraction

$$\iota_B : \Omega(X) \longrightarrow \Omega(X)$$

is continuous, and if Y is another manifold (possibly with boundary) and if $\Psi : X \rightarrow Y$ is a smooth map, then the pullback map

$$\Psi^* : \Omega(Y) \longrightarrow \Omega(X)$$

is continuous [17].

For every continuous seminorm ε on $\Omega(N)$ we get a seminorm $\varepsilon^{\mathbb{T}}$ on $\Omega_{\mathbb{T}}(N \times \mathbb{T})$ by setting

$$\varepsilon^{\mathbb{T}}(\alpha + \vartheta_{\mathbb{T}} \wedge \beta) := \varepsilon(\alpha) + \varepsilon(\beta)$$

In view of the formula $d_{\mathbb{T}}$, the space $\Omega_{\mathbb{T}}(N \times \mathbb{T})$ then becomes a locally convex unital DGA (by remark 3.1) in terms of the $\varepsilon^{\mathbb{T}}$'s. As a consequence, we get the super complexes

$$\mathcal{C}^+(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \xrightarrow{b+B} \mathcal{C}^-(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \xrightarrow{b+B} \mathcal{C}^+(\Omega_{\mathbb{T}}(N \times \mathbb{T})), \quad (10)$$

$$\mathcal{N}^+(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \xrightarrow{b+B} \mathcal{N}^-(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \xrightarrow{b+B} \mathcal{N}^+(\Omega_{\mathbb{T}}(N \times \mathbb{T})), \quad (11)$$

$$\mathcal{C}_{\varepsilon}^+(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \xrightarrow{b+B} \mathcal{C}_{\varepsilon}^-(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \xrightarrow{b+B} \mathcal{C}_{\varepsilon}^+(\Omega_{\mathbb{T}}(N \times \mathbb{T})), \quad (12)$$

$$\mathcal{N}_{\varepsilon}^+(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \xrightarrow{b+B} \mathcal{N}_{\varepsilon}^-(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \xrightarrow{b+B} \mathcal{N}_{\varepsilon}^+(\Omega_{\mathbb{T}}(N \times \mathbb{T})). \quad (13)$$

4 Equivariant Chen integrals

Let us consider a compact manifold N without boundary, and the space LN of smooth loops $\gamma : \mathbb{T} \rightarrow N$, where in the sequel we read \mathbb{T} as $\mathbb{T} = [0, 1]/\sim$. This becomes an infinite dimensional Fréchet manifold which is locally modelled on the Fréchet space $L\mathbb{R}^{\dim N}$ of smooth loops $\mathbb{T} \rightarrow \mathbb{R}^{\dim N}$. Then LN carries a natural smooth \mathbb{T} -action, given by rotating each loop, and the fixed point set of this action is precisely $N \subset LN$, embedded as constant loops. Given $\gamma \in LN$ the tangent space $T_{\gamma}LN$ is given by linear space of smooth vector fields on N along γ , that is,

$$T_{\gamma}(LN) = \{X \in C^{\infty}(\mathbb{T}, N) : X(t) \in T_{\gamma(t)}N \text{ for all } t \in \mathbb{T}\},$$

and the generator of the \mathbb{T} -action on LN is the vector field $\gamma \mapsto \dot{\gamma}$ on LN . Let ι denote the contraction with respect to the latter vector field. In the sequel, we understand

$$\Omega(LN) := \bigoplus_{k=0}^{\infty} \Omega^k(LN).$$

For fixed $s \in \mathbb{T}$ one has the diffeomorphism

$$\phi_s : LN \longrightarrow LN, \quad \gamma \longmapsto \gamma(s + \cdot)$$

induced by the \mathbb{T} -action, and one gets an induced operator

$$P : \Omega(LN) \longrightarrow \Omega(LN), \quad \text{defined on } \Omega^k(LN) \text{ by } P\alpha := \int_0^1 \phi_s^* \iota \alpha \, ds.$$

Then P becomes a degree -1 derivation. In addition, there is the usual exterior derivative

$$d : \Omega(LN) \longrightarrow \Omega(LN),$$

a degree +1 derivation. Taking only odd/even degree forms, one gets the super-structure $\Omega = \Omega^+(LN) \oplus \Omega^-(LN)$, and we get the super complex

$$\Omega^+(LN) \xrightarrow{d+P} \Omega^-(LN) \xrightarrow{d+P} \Omega^+(LN), \quad (14)$$

called the *equivariant de Rham complex of LN*. This complex does not carry much information, as the differential forms of interest, like the Bismut-Chern character below, are actually elements of

$$\prod_{k=0}^{\infty} \Omega^k(LN), \quad \text{rather than} \quad \Omega(LN) = \bigoplus_{k=0}^{\infty} \Omega^k(LN).$$

Thus we are going to 'complete' $\Omega(LN)$ in some way. To this end, following Chen's approach [6] of constructing a smooth structure on LN in terms of plots, we consider smooth maps $f : X \rightarrow LN$, where X is a finite dimensional manifold (without boundary). Given a continuous seminorm ε on $\Omega(X)$ we get an induced seminorm

$$\varepsilon_f(\omega) := \varepsilon(f^*\omega) \quad \text{on } \Omega(LN).$$

The locally convex topology induced by the ε_f 's is Hausdorff and we define $\widehat{\Omega}(LN)$ to be the completion of $\Omega(LN)$ with respect to this locally convex topology. The maps d, P and the grading operator become continuous maps $\Omega(LN) \rightarrow \Omega(LN)$: indeed, the continuity of the grading map is trivial. The continuity of d follows from

$$\varepsilon_f(d\omega) = \varepsilon(d[f^*\omega]) \leq \varepsilon'(f^*\omega) = \varepsilon'_f(\omega)$$

for some continuous seminorm ε' on $\Omega(X)$, where we have used the continuity of $d : \Omega(X) \rightarrow \Omega(X)$. Finally, the continuity of P follows easily from the continuity of ι , which in turn follows from writing

$$\varepsilon_f(\iota\omega) = \varepsilon(f^*[\iota\omega]) = \varepsilon(r^*\iota_{\partial_{\mathbb{T}}}\hat{f}^*j^*[\omega]) \leq \varepsilon'_{j \circ \hat{f}}(\omega)$$

for some continuous seminorm ε' on $\Omega(X \times \mathbb{T})$, where

$$r : X \longrightarrow X \times \mathbb{T}, \quad j : N \longrightarrow LN$$

are the canonical embeddings, and

$$\hat{f} : X \times \mathbb{T} \longrightarrow N$$

the map induced by $f : X \rightarrow LN$, and where we have used Remark 3.1 (the continuity of $r^*\iota_{\partial_{\mathbb{T}}}$, which implies the existence of ε').

We end up with the super complex

$$\widehat{\Omega}^+(LN) \xrightarrow{d+P} \widehat{\Omega}^-(LN) \xrightarrow{d+P} \widehat{\Omega}^+(LN), \quad (15)$$

called the *completed equivariant de Rham complex of LN*. The corresponding homology groups are denoted by $\widehat{H}_{\mathbb{T}}^{\pm}(LN)$.

Given $t \in \mathbb{T}$ and $\alpha \in \Omega^k(N)$ one denotes with $\alpha(t) \in \Omega^k(LN)$ the form obtained by pulling α back with respect to the evaluation map $\gamma \mapsto \gamma(t)$. With this notation at hand, one has the *equivariant Chen integral map*

$$\rho : \mathcal{C}(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \longrightarrow \Omega(LN),$$

which is defined by

$$\begin{aligned} & \rho \langle (\alpha_0 + \vartheta_{\mathbb{T}} \wedge \beta_0) \otimes \cdots \otimes (\alpha_n + \vartheta_{\mathbb{T}} \wedge \beta_n) \rangle \\ & := \int_0^1 ds \phi_s^* \int_{\Delta_n} \alpha_0(0) \wedge (\iota \alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota \alpha_n(t_n) - \beta_n(t_n)) dt_1 \cdots dt_n, \end{aligned}$$

where

$$\Delta_n = \{0 \leq t_1 \leq \cdots \leq t_n \leq 1\} \subset \mathbb{R}^n$$

denotes the standard n -simplex. We will also write

$$\begin{aligned} & \rho \langle (\alpha_0 + \vartheta_{\mathbb{T}} \wedge \beta_0) \otimes \cdots \otimes (\alpha_n + \vartheta_{\mathbb{T}} \wedge \beta_n) \rangle \\ & = \int_0^1 ds \phi_s^* \bar{\rho} \langle (\alpha_0 + \vartheta_{\mathbb{T}} \wedge \beta_0) \otimes \cdots \otimes (\alpha_n + \vartheta_{\mathbb{T}} \wedge \beta_n) \rangle. \end{aligned}$$

We collect the essential properties of ρ in the following proposition:

Proposition 4.1. *The map ρ is a continuous morphism of super complexes*

$$\rho : \mathcal{C}(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \longrightarrow \Omega(LN), \quad (16)$$

which in turn descends to a continuous map of super complexes

$$\rho : \mathcal{N}(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \longrightarrow \Omega(LN). \quad (17)$$

In particular, by density, we obtain the continuous maps of super complexes

$$\rho : \mathcal{C}_\epsilon(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \longrightarrow \widehat{\Omega}(LN), \quad \rho : \mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \longrightarrow \widehat{\Omega}(LN).$$

Proof. i) The fact that (16) is a map of superspaces follows easily from observing that

$$\begin{aligned} \mathcal{C}^+(\Omega_{\mathbb{T}}(N \times \mathbb{T})) &= \bigoplus_{j=0}^{\infty} \mathcal{C}^{2j}(\Omega_{\mathbb{T}}(N \times \mathbb{T})), \\ \mathcal{C}^-(\Omega_{\mathbb{T}}(N \times \mathbb{T})) &= \bigoplus_{j=0}^{\infty} \mathcal{C}^{2j+1}(\Omega_{\mathbb{T}}(N \times \mathbb{T})), \end{aligned}$$

where

$$\begin{aligned} & \mathcal{C}^k(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \\ &= \bigoplus_{r=0}^{\infty} \bigoplus_{l_0 + \cdots + l_r = k+r} \Omega_{\mathbb{T}}^{l_0}(N \times \mathbb{T}) \otimes \underline{\Omega_{\mathbb{T}}^{l_1}(N \times \mathbb{T})} \otimes \cdots \otimes \underline{\Omega_{\mathbb{T}}^{l_r}(N \times \mathbb{T})}, \end{aligned}$$

and that ρ maps $\mathcal{C}^k(\Omega_{\mathbb{T}}(N \times \mathbb{T})) \rightarrow \Omega^k(LN)$.

ii) Next we show that $\rho(b + B) = (d + P)\rho$. Setting $\omega_j = \alpha_j + \vartheta_{\mathbb{T}} \wedge \beta_j$, we first notice

$$\begin{aligned}
\tilde{\rho} b \langle \omega_0 \otimes \cdots \otimes \omega_n \rangle &= \tilde{\rho} \langle d_{\mathbb{T}} \omega_0 \otimes \cdots \otimes \omega_{j-1} \otimes \omega_j \otimes \omega_{j+1} \otimes \cdots \otimes \omega_n \rangle \\
&\quad - \tilde{\rho} \left\langle \sum_{j=1}^n (-1)^{r_{j-1}} \omega_0 \otimes \cdots \otimes \omega_{j-1} \otimes d_{\mathbb{T}} \omega_j \otimes \omega_{j+1} \otimes \cdots \otimes \omega_n \right\rangle \\
&\quad - \tilde{\rho} \left\langle \sum_{j=0}^{n-1} (-1)^{r_j} \omega_0 \otimes \cdots \otimes \omega_{j-1} \otimes \omega_j \wedge \omega_{j+1} \otimes \omega_{j+2} \otimes \cdots \otimes \omega_n \right\rangle \\
&\quad + (-1)^{(j_n-1)r_{n-1}} \tilde{\rho} \langle \omega_n \wedge \omega_0 \otimes \omega_1 \otimes \cdots \otimes \omega_{n-1} \rangle. \tag{18}
\end{aligned}$$

The first two lines give

$$\begin{aligned}
&\int_{\Delta_n} (d\alpha_0(0) + \beta_0(0)) \wedge (\iota\alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota\alpha_n(t_n) - \beta_n(t_n)) d^n t \\
&- \sum_{j=1}^n (-1)^{r_{j-1}} \int_{\Delta_n} \alpha_0(0) (\iota\alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota\alpha_{j-1}(t_{j-1}) - \beta_{j-1}(t_{j-1})) \wedge \\
&\quad \wedge (\iota d\alpha_j(t_j) + \iota\beta_j(t_{j-1}) + d\beta_j(t_j)) \wedge (\iota\alpha_{j+1}(t_{j+1}) - \beta_{j+1}(t_{j+1})) \wedge \cdots \wedge (\iota\alpha_n(t_n) - \beta_n(t_n)) d^n t,
\end{aligned}$$

where $d^n t = dt_1 \cdots dt_n$. Using that

$$\Delta_n = \{(t_1, t_2, \dots, t_n) : 0 \leq t_1 \leq \dots \leq t_{j-1} \leq t_j \leq t_{j+1} \leq \dots \leq t_n\},$$

and that

$$\iota d\alpha_j(t_j) = \frac{d}{dt_j} \alpha_j(t_j) - d\alpha_j(t_j),$$

it can be rewritten as

$$\begin{aligned}
&\int_{\Delta_n} (d\alpha_0(0) + \beta_0(0)) \wedge (\iota\alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota\alpha_n(t_n) - \beta_n(t_n)) d^n t \\
&+ \sum_{j=1}^n (-1)^{r_{j-1}} \int_{\Delta_n} \alpha_0(0) (\iota\alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota\alpha_{j-1}(t_{j-1}) - \beta_{j-1}(t_{j-1})) \wedge \\
&\quad \wedge d(\iota\alpha_j(t_j) - \beta_j(t_j)) \wedge (\iota\alpha_{j+1}(t_{j+1}) - \beta_{j+1}(t_{j+1})) \wedge \cdots \wedge (\iota\alpha_n(t_n) - \beta_n(t_n)) d^n t \\
&- \sum_{j=1}^n (-1)^{r_{j-1}} \int_{\Delta_n} \alpha_0(0) (\iota\alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota\alpha_{j-1}(t_{j-1}) - \beta_{j-1}(t_{j-1})) \wedge \\
&\quad \wedge \frac{d}{dt_j} \alpha_j(t_j) \wedge (\iota\alpha_{j+1}(t_{j+1}) - \beta_{j+1}(t_{j+1})) \wedge \cdots \wedge (\iota\alpha_n(t_n) - \beta_n(t_n)) d^n t \\
&- \sum_{j=1}^n (-1)^{r_{j-1}} \int_{\Delta_n} \alpha_0(0) (\iota\alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota\alpha_{j-1}(t_{j-1}) - \beta_{j-1}(t_{j-1})) \wedge \\
&\quad \wedge \iota\beta_j(t_j) \wedge (\iota\alpha_{j+1}(t_{j+1}) - \beta_{j+1}(t_{j+1})) \wedge \cdots \wedge (\iota\alpha_n(t_n) - \beta_n(t_n)) d^n t.
\end{aligned}$$

The first two (three) lines give

$$d\tilde{\rho}\langle\omega_0\otimes\cdots\otimes\omega_n\rangle+\int_{\Delta_n}\beta_0(0)\wedge(\iota\alpha_1(t_1)-\beta_1(t_1))\wedge\cdots\wedge(\iota\alpha_n(t_n)-\beta_n(t_n))d^n t, \quad (19)$$

while the third (fourth and fifth) line can be integrated in t_j from t_{j-1} to t_{j+1} thus getting

$$\begin{aligned} & d\tilde{\rho}\langle\omega_0\otimes\cdots\otimes\omega_n\rangle+\int_{\Delta_n}\beta_0(0)\wedge(\iota\alpha_1(t_1)-\beta_1(t_1))\wedge\cdots\wedge(\iota\alpha_n(t_n)-\beta_n(t_n))d^n t \\ & -\sum_{j=1}^{n-1}(-1)^{r_{j-1}}\int_{\Delta_{n-1}}\alpha_0(0)\wedge(\iota\alpha_1(t_1)-\beta_1(t_1))\wedge\cdots\wedge(\iota\alpha_{j-1}(t_{j-1})-\beta_{j-1}(t_{j-1}))\wedge \\ & \quad \wedge\alpha_j(t_{j+1})\wedge(\iota\alpha_{j+1}(t_{j+1})-\beta_{j+1}(t_{j+1}))\wedge\cdots\wedge(\iota\alpha_n(t_n)-\beta_n(t_n))d^n t_j \\ & -(-1)^{r_{n-1}}\int_{\Delta_{n-1}}\alpha_0(0)\wedge(\iota\alpha_1(t_1)-\beta_1(t_1))\wedge\cdots\wedge(\iota\alpha_{n-1}(t_{n-1})-\beta_{n-1}(t_{n-1}))\wedge\alpha_n(1)d^n t_n \\ & +\sum_{j=2}^n(-1)^{r_{j-1}}\int_{\Delta_{n-1}}\alpha_0(0)\wedge(\iota\alpha_1(t_1)-\beta_1(t_1))\wedge\cdots\wedge(\iota\alpha_{j-1}(t_{j-1})-\beta_{j-1}(t_{j-1}))\wedge \\ & \quad \wedge\alpha_j(t_{j-1})\wedge(\iota\alpha_{j+1}(t_{j+1})-\beta_{j+1}(t_{j+1}))\wedge\cdots\wedge(\iota\alpha_n(t_n)-\beta_n(t_n))d^n t_j \\ & +(-1)^{r_0}\int_{\Delta_{n-1}}\alpha_0(0)\wedge\alpha_1(0)\wedge(\iota\alpha_2(t_2)-\beta_2(t_2))\wedge\cdots\wedge(\iota\alpha_n(t_n)-\beta_n(t_n))d^n t_1 \\ & -\sum_{j=1}^n(-1)^{r_{j-1}}\int_{\Delta_n}\alpha_0(0)\wedge(\iota\alpha_1(t_1)-\beta_1(t_1))\wedge\cdots\wedge(\iota\alpha_{j-1}(t_{j-1})-\beta_{j-1}(t_{j-1}))\wedge \\ & \quad \wedge\iota\beta_j(t_j)\wedge(\iota\alpha_{j+1}(t_{j+1})-\beta_{j+1}(t_{j+1}))\wedge\cdots\wedge(\iota\alpha_n(t_n)-\beta_n(t_n))d^n t, \quad (20) \end{aligned}$$

where $d^n t_j = dt_1\cdots dt_{j-1}dt_{j+1}\cdots dt_n$. If in the fourth sum of integrals we change the summation variable from j to $j+1$, then make the change of variable $t_j \rightarrow t_{j+1}$, and put it together with the second sum of integrals, after noting that $(-1)^{r_{j-1}}(-1)^{j_j} = -(-1)^{r_j}$, then summing the fourth and the second integrals we get

$$\begin{aligned} & -\sum_{j=1}^{n-1}(-1)^{r_{j-1}}\int_{\Delta_{n-1}}\alpha_0(0)\wedge(\iota\alpha_1(t_1)-\beta_1(t_1))\wedge\cdots\wedge(\iota\alpha_{j-1}(t_{j-1})-\beta_{j-1}(t_{j-1}))\wedge \\ & \quad \wedge[\alpha_j(t_{j+1})\wedge(\iota\alpha_{j+1}(t_{j+1})-\beta_{j+1}(t_{j+1}))]\wedge\cdots\wedge(\iota\alpha_n(t_n)-\beta_n(t_n))d^n t_j \\ & +\sum_{j=1}^{n-1}(-1)^{r_j}\int_{\Delta_{n-1}}\alpha_0(0)\wedge(\iota\alpha_1(t_1)-\beta_1(t_1))\wedge\cdots\wedge(\iota\alpha_{j-1}(t_{j-1})-\beta_{j-1}(t_{j-1}))\wedge \\ & \quad \wedge[(\iota\alpha_j(t_{j+1})-\beta_j(t_{j+1}))\wedge\alpha_{j+1}(t_{j+1})]\wedge\cdots\wedge(\iota\alpha_n(t_n)-\beta_n(t_n))d^n t_j \\ & =\sum_{j=1}^{n-1}(-1)^{r_j}\int_{\Delta_{n-1}}\alpha_0(0)\wedge(\iota\alpha_1(t_1)-\beta_1(t_1))\wedge\cdots\wedge(\iota\alpha_{j-1}(t_{j-1})-\beta_{j-1}(t_{j-1}))\wedge \\ & \quad \wedge\left[(\iota\alpha_j(t_{j+1})-\beta_j(t_{j+1}))\wedge\alpha_{j+1}(t_{j+1})+(-1)^{j_j-1}\alpha_j(t_{j+1})\wedge(\iota\alpha_{j+1}(t_{j+1})-\beta_{j+1}(t_{j+1}))\right]\wedge \\ & \quad \wedge\cdots\wedge(\iota\alpha_n(t_n)-\beta_n(t_n))d^n t_j \\ & =\sum_{j=1}^{n-1}(-1)^{r_j}\tilde{\rho}\langle\omega_0\otimes\cdots\otimes\omega_{j-1}\otimes\omega_j\wedge\omega_{j+1}\otimes\omega_{j+2}\otimes\cdots\otimes\omega_n\rangle, \end{aligned}$$

which including the fifth integral in (4) becomes

$$\tilde{\rho} \left\langle \sum_{j=0}^{n-1} (-1)^{r_j} \omega_0 \otimes \cdots \otimes \omega_{j-1} \otimes \omega_j \wedge \omega_{j+1} \otimes \omega_{j+2} \otimes \cdots \otimes \omega_n \right\rangle.$$

This cancels the second line of (18). After noting that $\alpha_n(1) = \alpha_n(0)$, we see that the third integral in (4) is just

$$-(-1)^{(j_n-1)r_{n-1}} \tilde{\rho} \langle \omega_n \wedge \omega_0 \otimes \omega_1 \otimes \cdots \otimes \omega_{n-1} \rangle,$$

which cancels the third line of (18). Thus, we get

$$\begin{aligned} \tilde{\rho} b \langle \omega_0 \otimes \cdots \otimes \omega_n \rangle &= d\tilde{\rho} \langle \omega_0 \otimes \cdots \otimes \omega_n \rangle \\ &+ \int_{\Delta_n} \beta_0(0) \wedge (\iota\alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota\alpha_n(t_n) - \beta_n(t_n)) d^n t \\ &- \sum_{j=1}^n (-1)^{r_{j-1}} \int_{\Delta_n} \alpha_0(0) \wedge (\iota\alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota\alpha_{j-1}(t_{j-1}) - \beta_{j-1}(t_{j-1})) \wedge \\ &\quad \wedge \iota\beta_j(t_j) \wedge (\iota\alpha_{j+1}(t_{j+1}) - \beta_{j+1}(t_{j+1})) \wedge \cdots \wedge (\iota\alpha_n(t_n) - \beta_n(t_n)) d^n t. \end{aligned} \quad (21)$$

Now, let us consider

$$\begin{aligned} P\tilde{\rho} \langle \omega_0 \otimes \cdots \otimes \omega_n \rangle &= \int_I ds \phi_s^* \iota \int_{\Delta_n} \alpha_0(0) \wedge (\iota\alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota\alpha_n(t_n) - \beta_n(t_n)) d^n t \\ &= \int_{I \times \Delta_n} \iota\alpha_0(s) \wedge (\iota\alpha_1(t_1 + s) - \beta_1(t_1 + s)) \wedge \cdots \wedge (\iota\alpha_n(t_n + s) - \beta_n(t_n + s)) d^n t ds \\ &- \sum_{j=1}^n (-1)^{r_{j-1}} \int_I ds \phi_s^* \int_{\Delta_n} \alpha_0(0) \wedge (\iota\alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota\alpha_{j-1}(t_{j-1}) - \beta_{j-1}(t_{j-1})) \wedge \\ &\quad \wedge \iota\beta_j(t_j) \wedge (\iota\alpha_{j+1}(t_{j+1}) - \beta_{j+1}(t_{j+1})) \wedge \cdots \wedge (\iota\alpha_n(t_n) - \beta_n(t_n)) d^n t, \end{aligned} \quad (22)$$

where now I must be identified with the circle \mathbb{T} , and where we used that

$$\iota(\iota\alpha_k(t_k) - \beta_k(t_k)) = -\iota\beta_k(t_k).$$

Now, for any given choice of $\bar{t} = (t_1, \dots, t_n)$ such that $0 \leq t_1 \leq \cdots \leq t_n \leq 1$, we can understand \mathbb{T} as the union of almost everywhere $n+1$ disjoint intervals defined by

$$I_j(\bar{t}) = \{s \in \mathbb{T} \mid t_{j-1} + s \leq 1, t_j + s - 1 \geq 0\}, \quad j = 1, \dots, n+1.$$

We see that

$$D_j = \{I_j(\bar{t}) \times \bar{t} \mid \bar{t} \in \Delta_n\}$$

is a $(n+1)$ -simplex for any given j , and

$$\bigcup_{j=1}^{n+1} D_j = I \times \Delta_n$$

while $D_j \cap D_k$ has zero measure if $j \neq k$. Therefore,

$$\begin{aligned}
& \int_{I \times \Delta_n} \iota \alpha_0(s) \wedge (\iota \alpha_1(t_1 + s) - \beta_1(t_1 + s)) \wedge \cdots \wedge (\iota \alpha_n(t_n + s) - \beta_n(t_n + s)) d^n t ds \\
&= \int_{I \times \Delta_n} \beta_0(s) \wedge (\iota \alpha_1(t_1 + s) - \beta_1(t_1 + s)) \wedge \cdots \wedge (\iota \alpha_n(t_n + s) - \beta_n(t_n + s)) d^n t ds \\
&+ \int_{I \times \Delta_n} (\iota \alpha_0(s) - \beta_0) \wedge (\iota \alpha_1(t_1 + s) - \beta_1(t_1 + s)) \wedge \cdots \wedge (\iota \alpha_n(t_n + s) - \beta_n(t_n + s)) d^n t ds \\
&= \int_I ds \phi_s^* \int_{\Delta_n} \beta_0(0) \wedge (\iota \alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota \alpha_n(t_n) - \beta_n(t_n)) d^n t ds \\
&+ \sum_{j=1}^{n+1} \int_{D_j} (\iota \alpha_0(s) - \beta_0(s)) \wedge (\iota \alpha_1(t_1 + s) - \beta_1(t_1 + s)) \wedge \cdots \wedge (\iota \alpha_n(t_n + s) - \beta_n(t_n + s)) d^n t ds.
\end{aligned}$$

Now, for any given j we introduce the variables

$$\begin{aligned}
\tau_k &= t_{j+k-1} + s - 1, \quad k = 1, \dots, n+1-j, \\
\tau_{n+2-j} &= s, \\
\tau_k &= t_{k+j-n-2} + s, \quad k = n+3-j, \dots, n+1 \quad (\text{if } j \geq 2).
\end{aligned}$$

In this coordinates we have

$$D_j = \{(\tau_1, \dots, \tau_{n+1}) | 0 \leq \tau_1 \leq \cdots \leq \tau_{n+1} \leq 1\} \equiv \Delta_{n+1}, \quad d^n t ds = d^{n+1} \tau,$$

and

$$\begin{aligned}
& (\iota \alpha_0(s) - \beta_0(s)) \wedge (\iota \alpha_1(t_1 + s) - \beta_1(t_1 + s)) \wedge \cdots \wedge (\iota \alpha_n(t_n + s) - \beta_n(t_n + s)) \\
&= (-1)^{r_{j-1}(r_n - r_j)} 1 \wedge (\iota \alpha_j(\tau_1) - \beta_j(\tau_1)) \wedge \cdots \wedge (\iota \alpha_n(\tau_{n-j+1}) - \beta_n(\tau_{n-j+1})) \wedge \\
&\quad \wedge (\iota \alpha_0(\tau_{n-j+2}) - \beta_0(\tau_{n-j+2})) \wedge \cdots \wedge (\iota \alpha_{j-1}(\tau_{n+1}) - \beta_{j-1}(\tau_{n+1})).
\end{aligned}$$

Integrating over $D_j = \Delta_{n+1}$ it becomes

$$\begin{aligned}
& \int_{D_j} (\iota \alpha_0(s) - \beta_0(s)) \wedge (\iota \alpha_1(t_1 + s) - \beta_1(t_1 + s)) \wedge \cdots \wedge (\iota \alpha_n(t_n + s) - \beta_n(t_n + s)) \\
&= \rho \left\langle (-1)^{r_{j-1}(r_n - r_j)} 1 \otimes \omega_j \otimes \cdots \otimes \omega_n \otimes \omega_0 \otimes \cdots \otimes \omega_{j-1} \right\rangle,
\end{aligned}$$

and after summation over j we finally get

$$\begin{aligned}
P\tilde{\rho} \langle \omega_0 \otimes \cdots \otimes \omega_n \rangle &= \tilde{\rho} B \langle \omega_0 \otimes \cdots \otimes \omega_n \rangle \\
&+ \int_I ds \phi_s^* \int_{\Delta_n} \beta_0(0) \wedge (\iota \alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota \alpha_n(t_n) - \beta_n(t_n)) d^n t ds \\
&- \sum_{j=1}^n (-1)^{r_{j-1}} \int_I ds \phi_s^* \int_{\Delta_n} \alpha_0(0) \wedge (\iota \alpha_1(t_1) - \beta_1(t_1)) \wedge \cdots \wedge (\iota \alpha_{j-1}(t_{j-1}) - \beta_{j-1}(t_{j-1})) \wedge \\
&\quad \wedge \iota \beta_j(t_j) \wedge (\iota \alpha_{j+1}(t_{j+1}) - \beta_{j+1}(t_{j+1})) \wedge \cdots \wedge (\iota \alpha_n(t_n) - \beta_n(t_n)) d^n t.
\end{aligned}$$

Notice that the second and third lines here are the means over \mathbb{T} of the corresponding terms in (21). After taking the mean of both expressions and subtracting each other, we finally get $\rho(b + B) = (d + P)\rho$ as desired.

iii) We now prove that $\tilde{\rho}$ vanishes on $\mathcal{D}(\Omega_{\mathbb{T}}(N \times \mathbb{T}))$. This implies that ρ vanishes on $\mathcal{D}(\Omega_{\mathbb{T}}(N \times \mathbb{T}))$, too. For elements of the form (4) the assertion immediately follows from the fact that $\iota f(t) = 0$, as $f(t)$ is a zero form. So, let us consider an element of the form (5). Since (recall that f is constant over \mathbb{T})

$$\iota df(t) = \frac{d}{dt} f(t),$$

and $df = d_{\mathbb{T}}f$, we can write

$$\begin{aligned} & \tilde{\rho}(\langle \omega_0 \otimes \cdots \otimes \omega_{r-1} f \otimes \omega_{r+1} \otimes \cdots \otimes \omega_n \rangle + \langle \omega_0 \otimes \cdots \otimes \omega_{r-1} \otimes df \otimes \omega_{r+1} \otimes \cdots \otimes \omega_n \rangle \\ & \quad - \langle \omega_0 \otimes \cdots \otimes \omega_{r-1} \otimes f \omega_{r+1} \otimes \cdots \otimes \omega_n \rangle) \\ &= \int_{\Delta_{n-1}} \alpha_0(0) \wedge \cdots \wedge (\iota \alpha_{r-1}(t_{r-1}) f(t_{r-1}) - \beta_{r-1}(t_{r-1}) f(t_{r-1})) \wedge (\iota \alpha_{r+1}(t_{r+1}) - \beta_{r+1}(t_{r+1})) \wedge \\ & \quad \wedge \cdots \wedge (\iota \alpha_n(t_n) - \beta_n(t_n)) d^n t_r \\ & \quad - \int_{\Delta_{n-1}} \alpha_0(0) \wedge \cdots \wedge (\iota \alpha_{r-1}(t_{r-1}) - \beta_{r-1}(t_{r-1})) \wedge (f(t_{r+1}) \iota \alpha_{r+1}(t_{r+1}) - f(t_{r+1}) \beta_{r+1}(t_{r+1})) \wedge \\ & \quad \wedge \cdots \wedge (\iota \alpha_n(t_n) - \beta_n(t_n)) d^n t_r \\ & \quad + \int_{\Delta_n} \alpha_0(0) \wedge \cdots \wedge (\iota \alpha_{r-1}(t_{r-1}) - \beta_{r-1}(t_{r-1})) \wedge \frac{d}{dt_r} f(t_r) \wedge (\iota \alpha_r(t_r) - \beta_r(t_r)) \wedge \\ & \quad \wedge \cdots \wedge (\iota \alpha_n(t_n) - \beta_n(t_n)) d^n t. \end{aligned}$$

After integrating t_r from t_{r-1} to t_{r+1} in the last term, we get exactly zero.

v) It remains to check the continuity of (16), which easily follow from the continuity of $\tilde{\rho}$. To see the latter, let X be a smooth manifold (without boundary), let ε be a continuous seminorm on $\Omega(X)$, and let $f : X \rightarrow LN$ be smooth. For $s \in \mathbb{T}$ let r_s denote the embedding

$$X \longrightarrow X \times \mathbb{T}, \quad x \longmapsto (x, s).$$

Then we have

$$\begin{aligned} & \varepsilon_f(\tilde{\rho}(\langle \alpha_0 + \vartheta_{\mathbb{T}} \wedge \beta_0 \rangle \otimes \cdots \otimes \langle \alpha_n + \vartheta_{\mathbb{T}} \wedge \beta_n \rangle)) \\ & \leq \int_{\Delta_n} \varepsilon(f^*[\alpha_0(0)]) \prod_{i=1}^n \varepsilon(f^*[\iota \alpha_i(t_i) - \beta_i(t_i)]) dt_1 \cdots dt_n \\ & = \int_{\Delta_n} \varepsilon(r_0^* \hat{f}^* \alpha_0) \prod_{i=1}^n \varepsilon(r_{t_i}^* \iota_{\partial_r} \hat{f}^* \alpha_i - r_{t_i}^* \hat{f}^* \beta_i) dt_1 \cdots dt_n \\ & \leq \int_{\Delta_n} \varepsilon(r_0^* \hat{f}^* \alpha_0) \prod_{i=1}^n \left(\varepsilon(r_{t_i}^* \iota_{\partial_r} \hat{f}^* \alpha_i) + \varepsilon(r_{t_i}^* \hat{f}^* \beta_i) \right) dt_1 \cdots dt_n \\ & \leq \int_{\Delta_n} \tilde{\varepsilon}(\alpha_0) \prod_{i=1}^n \left(\tilde{\varepsilon}(\alpha_i) + \tilde{\varepsilon}(\beta_i) \right) dt_1 \cdots dt_n \\ & \leq \frac{1}{n!} \prod_{i=0}^n \left(\tilde{\varepsilon}(\alpha_i) + \tilde{\varepsilon}(\beta_i) \right) = \frac{1}{n!} \tilde{\varepsilon}_n^{\mathbb{T}} \left(\langle \alpha_0 + \vartheta_{\mathbb{T}} \wedge \beta_0 \rangle \otimes \cdots \otimes \langle \alpha_n + \vartheta_{\mathbb{T}} \wedge \beta_n \rangle \right), \end{aligned}$$

for some continuous seminorm $\tilde{\varepsilon}$ on $\Omega(N)$. This estimate shows the continuity of $\tilde{\rho}$ and completes the proof. ■

5 Construction of cycles in $\mathcal{N}_\epsilon^-(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$ and the induced cycles in $\widehat{\Omega}^-(LM)$

Let now M be a compact manifold (possibly with boundary). Given $g \in C^\infty(M, U(l \times l; \mathbb{C}))$ our aim is to construct a canonically given element

$$\text{Ch}^-(g) \in \mathcal{C}_\epsilon^-(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$$

with $(b+B)\text{Ch}^-(g) = 0$ in the Chen normalized complex. To this end, let $I := [0, 1]$ and denote the canonical vector field on I with ∂_I . We denote the canonical Maurer-Cartan form on $U(l \times l; \mathbb{C})$ by

$$\omega \in \Omega^1(U(l \times l; \mathbb{C}), \text{Mat}(l \times l; \mathbb{C})).$$

Then for all $s \in I$ we can form the covariant derivative $d + s\omega$ on the trivial vector bundle $U(l \times l; \mathbb{C}) \times \mathbb{C}^l \rightarrow U(l \times l; \mathbb{C})$. Let

$$A^s \in \Omega^1(U(l \times l; \mathbb{C}), \text{Mat}(l \times l; \mathbb{C})), \quad R^s \in \Omega^2(U(l \times l; \mathbb{C}), \text{Mat}(l \times l; \mathbb{C}))$$

denote the connection 1-form of $d + s\omega$ and the curvature of $d + s\omega$, respectively, and

$$\mathcal{A}^s := A^s - \vartheta_{\mathbb{T}} \wedge R^s \in \Omega_{\mathbb{T}}(U(l \times l; \mathbb{C}) \times \mathbb{T}, \text{Mat}(l \times l; \mathbb{C})).$$

We set

$$A^s(g) := g^* A^s, \quad R_g^s := g^* R^s, \quad \omega_g := g^* \omega,$$

so that $A^s(g) = s\omega_g$ and by the Maurer-Cartan equation $R_g^s = (s/2)\omega_g^2$. Then we can define

$$\mathcal{A}^s(g) := A_g^s - \vartheta_{\mathbb{T}} \wedge R_g^s \in \Omega_{\mathbb{T}}(M \times \mathbb{T}, \text{Mat}(l \times l; \mathbb{C})).$$

By varying s , the forms $\mathcal{A}^s(g)$ induce a form

$$\mathcal{A}(g) \in \Omega_{\mathbb{T}}(M \times I \times \mathbb{T}, \text{Mat}(l \times l; \mathbb{C}))$$

and we set

$$\mathcal{B}(g) := \iota_{\partial_I} \mathcal{A}(g) \in \Omega_{\mathbb{T}}(M \times I \times \mathbb{T}, \text{Mat}(l \times l; \mathbb{C})).$$

Then we can define

$$\mathcal{B}^s(g) \in \Omega_{\mathbb{T}}(M \times \mathbb{T}, \text{Mat}(l \times l; \mathbb{C})),$$

to be the pullback of $\mathcal{B}(g)$ with respect to the embedding

$$M \times \mathbb{T} \longrightarrow M \times I \times \mathbb{T}, \quad (x, t) \longmapsto (x, s, t).$$

In fact, by a simple calculation one finds

$$\mathcal{A}^s(g) = s\omega_g + s(1-s)\vartheta_{\mathbb{T}} \wedge \omega_g^2, \quad \mathcal{B}^s(g) = -\vartheta_{\mathbb{T}} \wedge \omega_g, \quad (23)$$

so that $\mathcal{B}^s(g)$ actually does not depend on s . With these preparations, we can define an element

$$\text{Ch}^-(g) = (\text{Ch}_0^-(g), \text{Ch}_1^-(g), \dots) \in \mathcal{C}(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$$

by setting

$$\text{Ch}_n^-(g) := \text{Tr}_n \left[\int_0^1 1 \otimes \sum_{k=1}^n \mathcal{A}^s(g)^{\otimes(k-1)} \otimes \mathcal{B}^s(g) \otimes \mathcal{A}^s(g)^{\otimes(n-k)} ds \right],$$

where given linear spaces V_0, \dots, V_n , and $v^{(j)} \in \text{Mat}(l \times l; V_j)$, $j = 0, \dots, n$, the generalized trace is defined by

$$\text{Tr}_n[v^{(0)} \otimes \dots \otimes v^{(n)}] := \sum_{i_0, \dots, i_n=1, \dots, l} v_{i_0, i_1}^{(0)} \otimes v_{i_1, i_2}^{(1)} \otimes \dots \otimes v_{i_n, i_0}^{(n)}.$$

We refer the reader to the paper [15] by Simons and Sullivan, where a construction of the usual odd Chern character $\text{ch}^-(g) \in \Omega^-(M)$ (cf. formula (24) below) has been given that influenced our definition of $\text{Ch}^-(g)$.

Theorem 5.1. *Let M be a compact manifold, possibly with boundary.*

a) *One has*

$$\text{Ch}^-(g) \in \mathcal{C}_\epsilon^-(\Omega_{\mathbb{T}}(M \times \mathbb{T})), \quad \text{and } (b+B)\text{Ch}^-(g) = 0 \text{ in } \mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T})),$$

in particular, $\text{Ch}^-(g)$ induces a homology class

$$[\text{Ch}^-(g)] \in \text{HN}_\epsilon^-(\Omega_{\mathbb{T}}(M \times \mathbb{T})).$$

b) *The map*

$$\text{K}^{-1}(M) \longrightarrow \text{HN}_\epsilon^-(\Omega_{\mathbb{T}}(M \times \mathbb{T})), \quad [g] \longmapsto [\text{Ch}^-(g)]$$

is a well-defined group homomorphism.

Proof. a) It is easily seen that $\Gamma\text{Ch}^-(g) = -\text{Ch}^-(g)$. To show that

$$\text{Ch}^-(g) \in \mathcal{C}_\epsilon^-(\Omega_{\mathbb{T}}(M \times \mathbb{T})),$$

given a continuous seminorm ϵ on $\Omega_{\mathbb{T}}(M \times \mathbb{T})$ set

$$C_\epsilon := \sup_{s \in [0,1]} \max \left(\epsilon(1), \max_{i,j=1, \dots, l} \epsilon(\mathcal{A}^s(g)_{ij}), \max_{i,j=1, \dots, l} \epsilon(\mathcal{B}^s(g)_{ij}) \right).$$

It is then easily checked that

$$\kappa_\epsilon(\text{Ch}^-(g)) \leq \sum_{n=0}^{\infty} n \frac{(l^2 C_\epsilon)^n}{\sqrt{n!}} < \infty.$$

It remains to prove

$$(b + B)\text{Ch}^-(g) \in \mathcal{D}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T})).$$

In fact,

$$B\text{Ch}^-(g) \in \mathcal{D}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T})),$$

as every $\langle \text{Ch}_n^-(g) \rangle$ contains the 0-form 1 and so is of the form (4) with $f = 1$. It remains to show that

$$b\text{Ch}^-(g) \in \mathcal{D}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T})).$$

In order to see the latter, let us first notice that

$$(b\text{Ch}^-(g))_n = (b\langle \text{Ch}_n^-(g) \rangle)_n + (b\langle \text{Ch}_{n+1}^-(g) \rangle)_n.$$

Using (23) and the explicit definition of b , we get

$$\begin{aligned} & (b\langle \text{Ch}_n^-(g) \rangle)_n \\ &= -\text{Tr}_n \left[\int_0^1 1 \otimes \sum_{k=1}^n \sum_{l=0}^{k-2} \mathcal{A}^s(g)^{\otimes l} \otimes (-s^2 \omega_g^2) \otimes \mathcal{A}^s(g)^{\otimes (k-l-2)} \right. \\ & \quad \left. \otimes (-\vartheta_{\mathbb{T}} \wedge \omega_g) \otimes \mathcal{A}^s(g)^{\otimes (n-k)} ds \right] \\ &+ \text{Tr}_n \left[\int_0^1 1 \otimes \sum_{k=1}^n \sum_{l=0}^{n-k-1} \mathcal{A}^s(g)^{\otimes (k-1)} \otimes (-\vartheta_{\mathbb{T}} \wedge \omega_g) \right. \\ & \quad \left. \otimes \mathcal{A}^s(g)^{\otimes l} \otimes (-s^2 \omega_g^2) \otimes \mathcal{A}^s(g)^{\otimes (n-k-l-1)} ds \right] \\ &- \text{Tr}_n \left[\int_0^1 1 \otimes \sum_{k=1}^n \mathcal{A}^s(g)^{\otimes (k-1)} \otimes (\vartheta_{\mathbb{T}} \wedge \omega_g^2 + \omega_g) \otimes \mathcal{A}^s(g)^{\otimes (n-k)} ds \right], \end{aligned}$$

and

$$\begin{aligned} & (b\langle \text{Ch}_{n+1}^-(g) \rangle)_n \\ &= -\text{Tr}_n \left[\int_0^1 1 \otimes \sum_{k=1}^n \sum_{l=0}^{k-2} \mathcal{A}^s(g)^{\otimes l} \otimes (+s^2 \omega_g^2) \otimes \mathcal{A}^s(g)^{\otimes (k-l-2)} \right. \\ & \quad \left. \otimes (-\vartheta_{\mathbb{T}} \wedge \omega_g) \otimes \mathcal{A}^s(g)^{\otimes (n-k)} ds \right] \\ &+ \text{Tr}_n \left[\int_0^1 1 \otimes \sum_{k=1}^n \sum_{l=0}^{n-k-1} \mathcal{A}^s(g)^{\otimes (k-1)} \otimes (-\vartheta_{\mathbb{T}} \wedge \omega_g) \otimes \mathcal{A}^s(g)^{\otimes l} \right. \\ & \quad \left. \otimes (+s^2 \omega_g^2) \otimes \mathcal{A}^s(g)^{\otimes (n-k-l-1)} ds \right] \\ &- \text{Tr}_n \left[\int_0^1 1 \otimes \sum_{k=1}^n \mathcal{A}^s(g)^{\otimes (k-1)} \otimes (-2s\vartheta_{\mathbb{T}} \wedge \omega_g^2) \otimes \mathcal{A}^s(g)^{\otimes (n-k)} ds \right], \end{aligned}$$

whose sum is

$$\begin{aligned} & \mathrm{Tr}_n \left[\int_0^1 1 \otimes \sum_{k=1}^n \mathcal{A}^s(g)^{\otimes(k-1)} \otimes \left(\frac{d}{ds} \mathcal{A}^s(g) \right) \otimes \mathcal{A}^s(g)^{\otimes(n-k)} ds \right] \\ &= \mathrm{Tr}_n \left[\int_0^1 \frac{d}{ds} (1 \otimes \mathcal{A}^s(g)^{\otimes n}) ds \right] = \mathrm{Tr}_n [1 \otimes \mathcal{A}^1(g)^{\otimes n}] - \mathrm{Tr}_n [1 \otimes \mathcal{A}^0(g)^{\otimes n}]. \end{aligned}$$

Thus, we finally have

$$(b\mathrm{Ch}^-(g))_n = \mathrm{Tr}_n [1 \otimes \omega_g^{\otimes n}], \quad n = 1, 2, \dots$$

We now prove that

$$(\dots, \mathrm{Tr}_n [1 \otimes \omega_g^{\otimes n}], \dots) \in \mathcal{D}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T})).$$

To this end we have simply to employ the properties of the generalized trace. Indeed, for $n \geq 2$ we can write

$$\begin{aligned} \langle \mathrm{Tr}_n [1 \otimes \omega_g^{\otimes n}] \rangle &= \left\langle \mathrm{Tr}_n [1 \otimes \omega_g \otimes \omega_g \otimes \omega_g^{\otimes(n-2)}] \right\rangle \\ &= - \left\langle \mathrm{Tr}_n [1 \otimes dg^{-1} \otimes dg \otimes \omega_g^{\otimes(n-2)}] \right\rangle \\ &= - \left\langle \mathrm{Tr}_n [1 \otimes dg^{-1} \otimes dg \otimes \omega_g^{\otimes(n-2)}] \right\rangle \\ &\quad - \left\langle \mathrm{Tr}_{n-1} [g^{-1} \otimes dg \otimes \omega_g^{\otimes(n-2)}] \right\rangle \\ &\quad + \left\langle \mathrm{Tr}_{n-1} [1 \otimes g^{-1} dg \otimes \omega_g^{\otimes(n-2)}] \right\rangle, \end{aligned}$$

where the last two terms cancel each other because of the trace property, which is precisely of the form (5) for $f = g^{-1}$. Similarly, for $n = 1$ it is sufficient to notice that

$$\langle \mathrm{Tr}_1 [1 \otimes \omega_g] \rangle = \langle \mathrm{Tr}_1 [g^{-1} \otimes dg] \rangle,$$

which is of the form (4) with $f = g^{-1}$, completing the proof of $b\mathrm{Ch}^-(g) \in \mathcal{D}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$.

b) It suffices to prove the following two facts:

- i) If $g, h \in C^\infty(M, U(l \times l; \mathbb{C}))$, then one has $\mathrm{Ch}^-(g \oplus h) = \mathrm{Ch}^-(g) + \mathrm{Ch}^-(h)$.
- ii) If $g_0, g_1 \in C^\infty(M, U(l \times l; \mathbb{C}))$ are connected by a smooth homotopy

$$g \in C^\infty(M \times I, U(l \times l; \mathbb{C})),$$

then one has

$$\mathrm{Ch}^-(g_1) - \mathrm{Ch}^-(g_0) = (b + B)w \quad \text{in } \mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$$

for some $w \in \mathcal{C}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$.

Here, property i) is an immediate consequence of the properties of the generalized

trace Tr_n using the block diagonal form of $g \oplus h$.
To see ii), for any $t \in I$, we define the embedding

$$j_t : M \hookrightarrow M \times I, \quad x \longmapsto (x, t),$$

and $w = (w_0, w_1, \dots) \in \mathcal{C}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$ by setting

$$\begin{aligned} w_n := & -\text{Tr}_n \left[\int_0^1 \int_0^1 1 \otimes \sum_{k=1}^n \sum_{l=0}^{k-2} j_t^* \left(\mathcal{A}^s(g.)^{\otimes l} \otimes \iota_{\partial_I} \mathcal{A}^s(g.) \otimes \mathcal{A}^s(g.)^{\otimes (k-l-2)} \right. \right. \\ & \left. \left. \otimes \mathcal{B}^s(g.) \otimes \mathcal{A}^s(g.)^{\otimes (n-k)} \right) ds dt \right] \\ & + \text{Tr}_n \left[\int_0^1 \int_0^1 1 \otimes \sum_{k=1}^n \sum_{l=0}^{n-k-1} j_t^* \left(\mathcal{A}^s(g.)^{\otimes (k-1)} \otimes \mathcal{B}^s(g.) \otimes \mathcal{A}^s(g.)^{\otimes l} \otimes \iota_{\partial_I} \mathcal{A}^s(g.) \right. \right. \\ & \left. \left. \otimes \mathcal{A}^s(g.)^{\otimes (n-k-l-1)} \right) ds dt \right] \\ & - \text{Tr}_n \left[\int_0^1 \int_0^1 1 \otimes \sum_{k=1}^n j_t^* \left(\mathcal{A}^s(g.)^{\otimes (k-1)} \otimes \iota_{\partial_I} \mathcal{B}^s(g.) \otimes \mathcal{A}^s(g.)^{\otimes (n-k)} \right) ds dt \right]. \end{aligned}$$

The \mathcal{C}_ϵ growth conditions are easily checked for w . Then again it is clear that $Bw \in \mathcal{D}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$. On the other hand, by using the identity

$$dj_t^* \iota_{\partial_I} \mathcal{A}^s(g.) = -j_t^* \iota_{\partial_I} d\mathcal{A}^s(g.) + \frac{\partial}{\partial t} j_t^* \mathcal{A}^s(g.),$$

and similarly for \mathcal{B}^s , and the same computations as in part a) we get, as elements in the Chen normalized complex,

$$\begin{aligned} (bw + Bw)_n &= (bw)_n = (b \langle w_n \rangle)_n + (b \langle w_{n+1} \rangle)_n = \left(\left\langle \int_0^1 \frac{d}{dt} j_t^* \text{Ch}^-(g.) \right\rangle_n \right) \\ &= \text{Ch}_n^-(g_1) - \text{Ch}_n^-(g_0). \end{aligned}$$

This completes the proof. ■

If M has no boundary (so that LM is a well-defined Fréchet manifold), in view of $(d + P)\rho = \rho(b + B)$, we immediately get:

Corollary 5.2. *Assume M is a compact manifold without boundary. Then for all $g \in C^\infty(M, U(l \times l; \mathbb{C}))$ one has $(d + P)\rho(\text{Ch}^-(g)) = 0$ in $\mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(M \times \mathbb{T}))$, in particular, $\rho(\text{Ch}^-(g))$ induces a homology class in $\widehat{H}_{\mathbb{T}}^-(LM)$.*

Remark 5.3. There is an even version of $\text{Ch}^-(g)$ given as follows: If N is a manifold and $d + C$ is a connection on a trivial vector bundle over N , then with R_C the curvature of the connection 1-form C one defines

$$\text{Ch}^+(C) = (\text{Ch}_0^+(C), \text{Ch}_1^+(C), \dots) \in \mathcal{C}_\epsilon^+(\Omega_{\mathbb{T}}(N \times \mathbb{T}))$$

by

$$\text{Ch}_n^+(C) := \text{Tr}_n [1 \otimes (C - \vartheta_{\mathbb{T}} \wedge R_C)^{\otimes n}],$$

which by an analogous calculation as in the proof of Theorem 5.1 is seen to satisfy

$$(b + B)\text{Ch}^+(C) = 0 \quad \text{in } \mathcal{N}_\epsilon(\Omega_{\mathbb{T}}(N \times \mathbb{T})).$$

Then, there holds an even/odd periodicity, that is, one can obtain $\text{Ch}^-(g)$ from its even variant by a fiber integration: indeed, by varying $s \in I$ in

$$A^s(g) \in \Omega_{\mathbb{T}}(M, \text{Mat}(l \times l; \mathbb{C}))$$

we get a form

$$A(g) \in \Omega_{\mathbb{T}}(M \times I, \text{Mat}(l \times l; \mathbb{C}))$$

and can consider the fibration

$$\pi : M \times I \longrightarrow M.$$

Then, for the connection $d + \tilde{A}_g$ on the trivial vector bundle over $M \times I$, where $\tilde{A}_g := \pi^* A_g$, one has, using the definitions of $\mathcal{A}^s(g)$ and $\mathcal{B}^s(g)$ that

$$\text{Ch}^-(g) = \int_I \iota_{\partial I} \text{Ch}^+(\tilde{A}_g) = \pi_* \text{Ch}^+(\tilde{A}_g),$$

the integration along the fibers of π .

The *odd Chern character* $\text{ch}^-(g) \in \Omega^-(M)$ is the closed odd differential form defined by

$$\text{ch}^-(g) := \text{Tr} \left[\sum_{j=0}^{\infty} \frac{(-1)^j j!}{(2j+1)!} (g^{-1} dg)^{\wedge(2j+1)} \right], \quad (24)$$

and the *odd Bismut-Chern character* is the differential form

$$\text{Bch}^-(g) = (\text{Bch}_1^-(g), \text{Bch}_3^-(g), \dots) \in \widehat{\Omega}^-(LM)$$

defined by

$$\text{Bch}_{2n-1}^-(g) = \text{Tr} \left[\int_0^1 \int_{\{0 \leq t_1 \leq \dots \leq t_n \leq 1\}} \sum_{j=1}^n \bigwedge_{i=1}^{j-1} //_{t_i}^s(g) R_g^s(t_i) \bigwedge //_{t_j}^s(g) \dot{A}_g^s(t_j) \right. \\ \left. \bigwedge_{l=j+1}^n //_{t_l}^s(g) R_g^s(t_l) //_{t_1}^s(g) dt_1 \cdots dt_n ds \right],$$

where

$$\dot{A}_g^s = \frac{d}{ds} A_g^s = \omega_g \in \Omega^1(M, \text{Mat}(l \times l; \mathbb{C})),$$

and where $//^s(g)$ denotes the parallel transport with respect to the connection $d + s\omega_g$ on the trivial vector bundle over M .

Theorem 5.4. *Assume M is a compact Riemannian manifold, possibly with boundary, and let $g \in C^\infty(M, U(l \times l; \mathbb{C}))$. Then one has $\rho(\text{Ch}^-(g))|_M = \text{ch}^-(g)$, and if M has no boundary then $\text{Bch}^-(g) = \rho(\text{Ch}^-(g))$.*

Note that in view of Corollary 5.2, Theorem 5.4 provides a new proof of

$$(d + P)\text{Bch}^-(g) = 0$$

We refer the reader to [18] for a variant of this result.

Proof of Theorem 5.4. The formula $\rho(\text{Ch}^-(g))|_M = \text{ch}^-(g)$ is a simple consequence of the definitions, once one has noticed the formula

$$\rho \langle (\alpha_0 + \vartheta_{\mathbb{T}} \wedge \beta_0) \otimes \cdots \otimes (\alpha_n + \vartheta_{\mathbb{T}} \wedge \beta_n) \rangle |_M = \alpha_0 \wedge \cdots \wedge \alpha_n.$$

In order to see $\text{Bch}^-(g) = \rho(g)$, given $t, s \in I$ define

$$V^s(g, t) \in \widehat{\Omega}^-(LM, \text{Mat}(l \times l; \mathbb{C}))$$

by

$$\begin{aligned} V_{2n+1}^s(g, t) = & \int_{\{0 \leq t_1 \leq \dots \leq t_{n+1} \leq t\}} \sum_{j=1}^{n+1} \bigwedge_{i=1}^{j-1} //_{t_i}^s(g) R_g^s(t_i) \bigwedge //_{t_j}^s(g) \dot{A}_g^s(t_j) \\ & \times \bigwedge_{l=j+1}^{n+1} //_{t_l}^s(g) R_g^s(t_l) //_1^s(g) dt_1 \cdots dt_{n+1}, \end{aligned}$$

and the differential form

$$W^s(g, t) \in \widehat{\Omega}^-(LM, \text{Mat}(l \times l; \mathbb{C}))$$

by

$$\begin{aligned} W_{2n+1}^s(g, t) = & \sum_{k=n+1}^{\infty} \sum_{r, j_1, \dots, j_n=1, \text{pairwise distinct}}^k \\ & \times \int_{\{0 \leq t_1 \leq \dots \leq t_k \leq t\}} \iota A_g^s(t_1) \cdots R_g^s(t_{j_1}) \cdots \dot{A}_g^s(t_r) \cdots R_g^s(t_{j_n}) \cdots \iota A_g^s(t_k) dt_1 \cdots dt_k. \end{aligned}$$

Then obviously one has

$$\text{Bch}^-(g) = \text{Tr} \left[\int_0^1 V^s(g, t)|_{t=1} ds \right]$$

and it is easily checked from the definitions that

$$\rho(\text{Ch}^-(g)) = \text{Tr} \left[\int_0^1 W^s(g, t)|_{t=1} ds \right].$$

Thus it suffices to show that $W^s(g, t) = V^s(g, t)$ for all $t, s \in I$. To see this, the essential idea is to consider for every $t, s \in I$ the even form

$$X^s(g, t) = (X_0^s(g, t), X_2^s(g, t), \dots) \in \widehat{\Omega}^+(LM, \text{Mat}(l \times l; \mathbb{C}))$$

which is defined by

$$\begin{aligned} X_0^s(g, t) &= //_t^s(g), \\ \frac{d}{dt} X_{2n}^s(g, t) &= X_{2n}^s(g, t) \iota A_g^s(t) + X_{2n-2}^s(g, t) R_g^s(t), \\ X_{2n}^s(g, t)|_{t=0} &= 0 \quad \text{for all } n \geq 1, \end{aligned}$$

and the odd form

$$Y^s(g, t) = (Y_1^s(g, t), Y_3^s(g, t), \dots) \in \Omega^-(LM, \text{Mat}(l \times l; \mathbb{C}))$$

which is defined by

$$\begin{aligned} \frac{d}{dt} Y_1^s(g, t) &= Y_1^s(g, t) \iota A_g^s(t) + X_0^s(g, t) \dot{A}_g^s(t), \\ \frac{d}{dt} Y_{2n+1}^s(g, t) &= Y_{2n+1}^s(g, t) \iota A_g^s(t) + Y_{2n-1}^s(g, t) R_g^s(t) + X_{2n}^s(g, t) \dot{A}_g^s(t) \quad \forall n \geq 1, \\ Y_{2n+1}^s(g, t)|_{t=0} &= 0 \quad \text{for all } n. \end{aligned}$$

Noting that the sum that defines $W_{2n+1}^s(g, t)$ converges uniformly in t so that one can interchange d/dt with $\sum_{k=n+1}^{\infty}$, it is now easily checked that both $t \mapsto W^s(g, t)$ and $t \mapsto V^s(g, t)$ solve the IVP's which define $Y^s(g, t)$, so that

$$V^s(g, t) = W^s(g, t) = Y^s(g, t) \quad \text{for all } t, s \in I,$$

as was claimed. ■

Remark 5.5. If N is a compact manifold without boundary and given a connection $d + C$ over a trivial vector bundle over N , the *even Bismut-Chern character* is the differential form

$$\text{Bch}^+(C) = (\text{Bch}_0^+(C), \text{Bch}_2^+(C), \dots) \in \widehat{\Omega}^+(LN)$$

defined by

$$\text{Bch}_{2n}^+(C) = \text{Tr} \left[\int_{\{0 \leq t_1 \leq \dots \leq t_n \leq 1\}} \bigwedge_{i=1}^n //_{t_i}^C R_C(t_i) //_1^C dt_1 \cdots dt_n \right],$$

where R_C is again the curvature of $d + C$ and $//^C$ is the parallel transport with respect to $d + C$. Then one has another even/odd periodicity as in Remark 5.3: we can consider A_g^s as defining a connection 1-form \dot{A}_g over a trivial vector bundle over $M \times I$. However, since $M \times I$ is a manifold with boundary, it is convenient to embed it in a larger manifold, say

$$\chi : M \times I \hookrightarrow M \times J$$

where $J = (-1, 2)$. Therefore, we extend A_g^s to $s \in J$, consider it as defining a connection 1-form \tilde{A}_g over a trivial vector bundle over $M \times J$. The corresponding curvature

$$R_{\tilde{A}_g} \in \Omega^2(M \times J, \text{Mat}(l \times l; \mathbb{C}))$$

is given by varying $s \in J$ in

$$R_g^s + ds \wedge \dot{A}_g^s \in \Omega^2(M, \text{Mat}(l \times l; \mathbb{C})).$$

Since $\iota_{\partial J} R_{\tilde{A}_g} = \dot{A}_g^s$, after restricting to loops fibering over J , we immediately get that under integration along the fibers of

$$\pi : M \times I \longrightarrow M,$$

one has

$$\text{Bch}_{2n-1}^-(g) = \int_I \chi^* \iota_{\partial J} \text{Bch}_{2n}^+(\tilde{A}_g) = \pi_* \chi^* \text{Bch}_{2n}^+(\tilde{A}_g).$$

References

- [1] Atiyah, M. : *Circular symmetry and stationary-phase approximation*. Colloquium in honor of Laurent Schwartz, vol. 1, (Palaiseau, 1983). Asterisque, 1(131):43–59, 1985.
- [2] Bismut, J.-M.: *Index theorem and equivariant cohomology on the loop space*. Comm. Math. Phys. 98 (1985), no. 2, 213–237.
- [3] Bismut, J.-M.: *Duistermaat-Heckman formulas and index theory*. Geometric aspects of analysis and mechanics, Progr. Math., vol. 292, Birkhauser/Springer, New York, 2011, pp. 1–55.
- [4] Bismut, J.-M. & Freed, D.S.: *The analysis of elliptic families. I. Metrics and connections on determinant bundles*. Comm. Math. Phys., Volume 106, Number 1 (1986), 159–176.
- [5] Bismut, J.-M. & Freed, D.S.: *The analysis of elliptic families. II. Dirac operators, eta invariants, and the holonomy theorem*. Comm. Math. Phys., Volume 107, Number 1 (1986), 103–163.
- [6] Chen, K.-T.: *Iterated integrals of differential forms and loop space homology*. Ann. of Math. (2) 97 (1973), 217–246.
- [7] Connes, A.: *Entire cyclic cohomology of Banach algebras and characters of θ -summable Fredholm modules*. K-Theory 1, (1988), 519–548.
- [8] Getzler, E.: *The odd Chern character in cyclic homology and spectral flow*. Topology 32 (1993), no. 3, 489–507.

- [9] Getzler, E. *The Thom class of Mathai and Quillen and probability theory*. Stochastic Analysis and Applications (Lisbon 1989). Prog. Probab. 26, Birkhauser, 1991, 111–122.
- [10] Getzler, E. & Szenes, A.: *On the Chern character of theta-summable Fredholm modules*. J. Func. Anal. 84 (1989), 343–357.
- [11] Getzler, E.; Jones, John D. S.; Petrack, Scott: *Differential forms on loop spaces and the cyclic bar complex*. Topology 30 (1991), no. 3, 339–371.
- [12] Getzler, E.; Szenes, A.: *On the Chern character of theta-summable Fredholm modules*. J. Func. Anal. 84 (1989), 343–357.
- [13] Hanisch, F. & Ludewig, M.: *Supersymmetric Path Integrals I: Differential Forms on the Loop Space*. arXiv:1709.10027.
- [14] Shen, Shu: Private communication.
- [15] Simons, J. & Sullivan, D.: *Structured Vector Bundles define Differential K-Theory*. arXiv:0810.4935.
- [16] Tradler, Thomas; Wilson, Scott O.; Zeinalian, Mahmoud: *Equivariant holonomy for bundles and abelian gerbes*. Comm. Math. Phys. 315 (2012), no. 1, 39–108.
- [17] Waldmann, S.: *Geometric Wave Equations*. arXiv:1208.4706v1.
- [18] Wilson, S.O: *A loop group extension of the odd Chern character*. J. Geom. Phys. 102 (2016), 32–43.