

Sparse high-dimensional FFT based on rank-1 lattice sampling

Toni Volkmer

Faculty of Mathematics
Technische Universität Chemnitz

<http://www.tu-chemnitz.de/~tovo>



joint work with Daniel Potts, Manfred Tasche

supported by



Content

Introduction

Multivariate trigonometric polynomials

High-dim. sparse FFT via 1-dim. sparse FFT

Sparse dimension-incremental FFT

Non-periodic case

Summary

Introduction

- approximate high-dim. function $f: \mathbb{T}^d \simeq [0, 1)^d \rightarrow \mathbb{C}$ by multivariate trigonometric polynomial $p: \mathbb{T}^d \rightarrow \mathbb{C}$ with frequencies supported on $I \subset \mathbb{Z}^d$, $|I| < \infty$,


$$p(\mathbf{x}) := \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}, \quad \hat{p}_{\mathbf{k}} \in \mathbb{C}$$

Introduction

- approximate high-dim. function $f: \mathbb{T}^d \simeq [0, 1)^d \rightarrow \mathbb{C}$ by multivariate trigonometric polynomial $p: \mathbb{T}^d \rightarrow \mathbb{C}$ with frequencies supported on $I \subset \mathbb{Z}^d$, $|I| < \infty$,

$$p(\mathbf{x}) := \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}, \quad \hat{p}_{\mathbf{k}} \in \mathbb{C}$$

- arbitrary known** frequency index set $I \subset \mathbb{Z}^d$, $|I| < \infty$, rank-1 lattice nodes \mathbf{x}_j , $j = 0, \dots, M-1$
 - fast evaluation $p(\mathbf{x}_j)$, (e.g. [Li, Hickernell 03])
 - fast and exact reconstruction of $\hat{p}_{\mathbf{k}}$, $\mathbf{k} \in I$, from samples $p(\mathbf{x}_j)$, ([Kämmerer, Kunis, Potts 12] [Kämmerer 13])
 - approximate reconstruction of f by p from samples $f(\mathbf{x}_j)$




Introduction

- approximate high-dim. function $f: \mathbb{T}^d \simeq [0, 1)^d \rightarrow \mathbb{C}$ by multivariate trigonometric polynomial $p: \mathbb{T}^d \rightarrow \mathbb{C}$ with frequencies supported on $I \subset \mathbb{Z}^d$, $|I| < \infty$,

$$p(\mathbf{x}) := \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}, \quad \hat{p}_{\mathbf{k}} \in \mathbb{C}$$

- arbitrary known** frequency index set $I \subset \mathbb{Z}^d$, $|I| < \infty$, rank-1 lattice nodes \mathbf{x}_j , $j = 0, \dots, M-1$
 - fast evaluation $p(\mathbf{x}_j)$, (e.g. [Li, Hickernell 03])
 - fast and exact reconstruction of $\hat{p}_{\mathbf{k}}$, $\mathbf{k} \in I$, from samples $p(\mathbf{x}_j)$, ([Kämmerer, Kunis, Potts 12] [Kämmerer 13])
 - approximate reconstruction of f by p from samples $f(\mathbf{x}_j)$




- unknown** frequency index set I ?

Trigonometric polynomials - fast evaluation

- rank-1 lattice $\text{R1L}(z, M)$: $z \in \mathbb{N}_0^d, M \in \mathbb{N}$

$$x_j = \frac{j}{M}z \bmod \mathbf{1}; j = 0, \dots, M-1$$



Korobov 59
Maisonneuve 72
Sloan & Kachoyan 84,87,90
Temlyakov 86
Lyness 89
Sloan & Joe 94
Sloan & Reztsov 01
Li & Hickernell 03

Trigonometric polynomials - fast evaluation



- rank-1 lattice R1L(\mathbf{z}, M): $\mathbf{z} \in \mathbb{N}_0^d, M \in \mathbb{N}$

$$\mathbf{x}_j = \frac{j}{M} \mathbf{z} \bmod \mathbf{1}; j = 0, \dots, M-1$$


- multivariate high-dim. trigonometric polynomial $p(\mathbf{x}) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$

- reformulation

$$p(\mathbf{x}_j) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \frac{j \mathbf{k} \cdot \mathbf{z}}{M}} = \sum_{l=0}^{M-1} \underbrace{\left(\sum_{\substack{\mathbf{k} \in I \\ \mathbf{k} \cdot \mathbf{z} \equiv l \pmod{M}} \hat{p}_{\mathbf{k}} \right)}_{\hat{g}_l} e^{2\pi i \frac{j \mathbf{k} \cdot \mathbf{z}}{M}} = \sum_{l=0}^{M-1} \hat{g}_l e^{2\pi i \frac{j l}{M}}$$



$(\hat{p}_{\mathbf{k}})_{\mathbf{k} \in I}$



$(p(\mathbf{x}_j))_{j=0}^{M-1}$



Trigonometric polynomials - fast evaluation

- rank-1 lattice R1L(\mathbf{z}, M): $\mathbf{z} \in \mathbb{N}_0^d, M \in \mathbb{N}$

$$\mathbf{x}_j = \frac{j}{M} \mathbf{z} \bmod \mathbf{1}; j = 0, \dots, M-1$$

- multivariate high-dim. trigonometric polynomial $p(\mathbf{x}) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$
- reformulation




$$p(\mathbf{x}_j) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \frac{j \mathbf{k} \cdot \mathbf{z}}{M}} = \sum_{l=0}^{M-1} \underbrace{\left(\sum_{\substack{\mathbf{k} \in I \\ \mathbf{k} \cdot \mathbf{z} \equiv l \pmod{M}} \hat{p}_{\mathbf{k}} \right)}_{\hat{g}_l} e^{2\pi i \frac{j \mathbf{k} \cdot \mathbf{z}}{M}} = \sum_{l=0}^{M-1} \hat{g}_l e^{2\pi i \frac{j l}{M}}$$



$(\hat{p}_{\mathbf{k}})_{\mathbf{k} \in I}$

1-dim
 $\xrightarrow{\text{FFT}}$

$$\mathcal{O}(M \log M + d|I|)$$



$(p(\mathbf{x}_j))_{j=0}^{M-1}$


Trigonometric polynomials - fast reconstruction

- rank-1 lattice $\text{R1L}(z, M)$: $z \in \mathbb{N}_0^d, M \in \mathbb{N}$

$$\mathbf{x}_j := \frac{j}{M} z \bmod \mathbf{1}; \quad j = 0, \dots, M-1$$

- reconstruction of Fourier coefficients $\hat{p}_{\mathbf{k}}$ of multivariate trigonometric polynomial

$$p(\mathbf{x}) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$$



Trigonometric polynomials - fast reconstruction

- rank-1 lattice R1L(z, M): $z \in \mathbb{N}_0^d, M \in \mathbb{N}$

$$\mathbf{x}_j := \frac{j}{M} \mathbf{z} \bmod \mathbf{1}; j = 0, \dots, M-1$$



- reconstruction of Fourier coefficients $\hat{p}_{\mathbf{k}}$ of multivariate trigonometric polynomial

$$p(\mathbf{x}) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$$

⇒ **Definition** reconstructing R1L(z, M) for I :


$$\mathbf{k} \cdot \mathbf{z} \not\equiv \mathbf{k}' \cdot \mathbf{z} \pmod{M} \text{ for all } \mathbf{k}, \mathbf{k}' \in I, \mathbf{k} \neq \mathbf{k}'$$

- $|I| \leq M \leq |I|^2$, CBC construction algorithm (Kämmerer 2012)



1-dim
←
iFFT

$$\mathcal{O}(M \log M + d|I|)$$



Trigonometric polynomials - fast approximation



- reconstructing rank-1 lattice for I ,

$$\mathbf{x}_j := \frac{j}{M} \mathbf{z} \bmod \mathbf{1}; j = 0, \dots, M-1$$

- approximation of function $f: \mathbb{T}^d \rightarrow \mathbb{C}$ by multivariate trigonometric polynomial


$$p(\mathbf{x}) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$$

- [Kuo, Sloan, Woźniakowski 06] [Kämmerer, Potts, V. 15] [Byrenheid, Kämmerer, Ullrich, V. 16]



1-dim
←
iFFT

$$\mathcal{O}(M \log M + d|I|)$$



Unknown frequency index set

until now: fast reconstruction of $p(\mathbf{x}) := \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$ or
fast approximation of function $f(\mathbf{x}) \approx p(\mathbf{x})$

- given frequency index set I
- compute $\hat{p}_{\mathbf{k}}$ from samples along reconstructing rank-1 lattice

Unknown frequency index set

until now: fast reconstruction of $p(\mathbf{x}) := \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$ or
fast approximation of function $f(\mathbf{x}) \approx p(\mathbf{x})$

- given frequency index set I
- compute $\hat{p}_{\mathbf{k}}$ from samples along reconstructing rank-1 lattice

next: unknown I

- search for location I of largest Fourier coefficients of f or non-zero Fourier coefficients of p
(and compute Fourier coefficients $\hat{p}_{\mathbf{k}}, \mathbf{k} \in I$)

Unknown frequency index set

until now: fast reconstruction of $p(\mathbf{x}) := \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$ or
fast approximation of function $f(\mathbf{x}) \approx p(\mathbf{x})$

- given frequency index set I
- compute $\hat{p}_{\mathbf{k}}$ from samples along reconstructing rank-1 lattice


next: unknown I

- search for location I of largest Fourier coefficients of f or non-zero Fourier coefficients of p
(and compute Fourier coefficients $\hat{p}_{\mathbf{k}}, \mathbf{k} \in I$)
- search domain: (possibly) large index set $\Gamma \subset \mathbb{Z}^d$, e.g., full grid $\hat{G}_N^d := \{\mathbf{k} \in \mathbb{Z}^d : \|\mathbf{k}\|_{\infty} \leq N\}$, ($|\hat{G}_{64}^{10}| \approx 1.28 \cdot 10^{21}$)


High-dim. sparse FFT via 1-dim. sparse FFT

General idea: use rank-1 lattice and 1-dim. sparse FFT

- Interpret $p(\mathbf{x}_j) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i(\mathbf{k} \cdot \mathbf{z}) \frac{j}{M}} =: \tilde{p}(\frac{j}{M})$
as evaluation of 1-dim. trigonometric polynomial $\tilde{p}: \mathbb{T} \rightarrow \mathbb{C}$
with frequencies $\mathbf{k} \cdot \mathbf{z} \bmod M$ at nodes j/M



1-dim
↔
FFT




High-dim. sparse FFT via 1-dim. sparse FFT

General idea: use rank-1 lattice and 1-dim. sparse FFT

- Interpret $p(\mathbf{x}_j) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i(\mathbf{k} \cdot \mathbf{z}) \frac{j}{M}} =: \tilde{p}(\frac{j}{M})$
as evaluation of 1-dim. trigonometric polynomial $\tilde{p}: \mathbb{T} \rightarrow \mathbb{C}$
with frequencies $\mathbf{k} \cdot \mathbf{z} \bmod M$ at nodes j/M

\Rightarrow reconstructing rank-1 lattice for $\Gamma \supset I$ gives injective mapping
of multi-dim. frequencies to 1-dim. frequencies,
 $\Gamma \subset \mathbb{Z}^d \rightarrow \{0, 1, \dots, M-1\}, \mathbf{k} \mapsto \mathbf{k} \cdot \mathbf{z} \bmod M$




High-dim. sparse FFT via 1-dim. sparse FFT

General idea: use rank-1 lattice and 1-dim. sparse FFT

- Interpret $p(\mathbf{x}_j) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i(\mathbf{k} \cdot \mathbf{z}) \frac{j}{M}} =: \tilde{p}(\frac{j}{M})$ as evaluation of 1-dim. trigonometric polynomial $\tilde{p}: \mathbb{T} \rightarrow \mathbb{C}$ with frequencies $\mathbf{k} \cdot \mathbf{z} \bmod M$ at nodes j/M

\Rightarrow reconstructing rank-1 lattice for $\Gamma \supset I$ gives injective mapping of multi-dim. frequencies to 1-dim. frequencies,
 $\Gamma \subset \mathbb{Z}^d \rightarrow \{0, 1, \dots, M-1\}, \mathbf{k} \mapsto \mathbf{k} \cdot \mathbf{z} \bmod M$

\Rightarrow 1-dim. sparse FFT methods can be applied



High-dim. sparse FFT via 1-dim. sparse FFT

usage of 1-dim. sparse FFT methods based on e.g.

	samples	computational costs
compressed sensing	$\mathcal{O}(I \log^4(M) \log(1/\eta))$ [Rauhut 07] [Kunis, Rauhut 08] [Gröchenig, Pötscher, Rauhut 10] [Foucart, Rauhut 13] ...	$\mathcal{O}(M I \log^4(M) \log(1/\eta))$
filters	$\mathcal{O}(I (\log M)(\log(M/ I)))$ [Hassanieh, Indyk, Katabi, Price 12] ...	$\mathcal{O}(I (\log M)(\log(M/ I)))$
C.R.T.	$\mathcal{O}(I \log^4(M))$ [Iwen 10] [Iwen 13]	$\mathcal{O}(I \log^4(M))$
shifted sampling	$\mathcal{O}(I \log(M/ I))$ (on average) [Christlieb, Lawlor, Wang 15]	$\mathcal{O}(I \log(M/ I))$
ESPRIT	$\mathcal{O}(I)$ (deterministic) [Roy, Kailath 89]	$\mathcal{O}(I ^3)$
ESPRIT + shifted samp.	$\mathcal{O}(I)$ (1 iteration) [Potts, Tasche, V. 16]	$\mathcal{O}(I ^{5/3})$

problems: constants?, noise / stability, “whp”, implementation


High-dim. sparse FFT via 1-dim. sparse FFT

usage of 1-dim. sparse FFT methods based on e.g.



	samples	computational costs
compressed sensing	$\mathcal{O}(I \log^4(M) \log(1/\eta))$ [Rauhut 07] [Kunis, Rauhut 08] [Gröchenig, Pötscher, Rauhut 10] [Foucart, Rauhut 13] ...	$\mathcal{O}(M I \log^4(M) \log(1/\eta))$
filters	$\mathcal{O}(I (\log M)(\log(M/ I)))$ [Hassanieh, Indyk, Katabi, Price 12] ...	$\mathcal{O}(I (\log M)(\log(M/ I)))$
C.R.T.	$\mathcal{O}(I \log^4(M))$ [Iwen 10] [Iwen 13]	$\mathcal{O}(I \log^4(M))$
shifted sampling	$\mathcal{O}(I \log(M/ I))$ (on average) [Christlieb, Lawlor, Wang 15]	$\mathcal{O}(I \log(M/ I))$
ESPRIT	$\mathcal{O}(I)$ (deterministic) [Roy, Kailath 89]	$\mathcal{O}(I ^3)$
ESPRIT + shifted samp.	$\mathcal{O}(I)$ (1 iteration) [Potts, Tasche, V. 16]	$\mathcal{O}(I ^{5/3})$

problems: constants?, noise / stability, “whp”, implementation
next: different approach based on CBC idea



Sparse dimension-incremental FFT - method




Sparse dimension-incremental FFT - method




Sparse dimension-incremental FFT - method



construct
→
sampling set




Sparse dimension-incremental FFT - method




$$\hat{p}_{k_1} := \frac{1}{17} \sum_{\ell=0}^{16} p \left(\begin{pmatrix} \ell/17 \\ x'_2 \\ x'_3 \end{pmatrix} \right) e^{-2\pi i \frac{\ell k_1}{17}}$$


$$k_1 = -8, \dots, 8$$



1-dim
←
iFFT




Sparse dimension-incremental FFT - method




$$\begin{aligned} \hat{p}_{k_1} &:= \frac{1}{17} \sum_{\ell=0}^{16} p \left(\begin{pmatrix} \ell/17 \\ x'_2 \\ x'_3 \end{pmatrix} \right) e^{-2\pi i \frac{\ell k_1}{17}} \\ &= \sum_{\substack{(h_2, h_3) \in \{-8, \dots, 8\}^2 \\ (k_1, h_2, h_3)^\top \in \text{supp } \hat{p}}} \hat{p} \begin{pmatrix} k_1 \\ h_2 \\ h_3 \end{pmatrix} e^{2\pi i (h_2 x'_2 + h_3 x'_3)}, \end{aligned}$$



$$k_1 = -8, \dots, 8$$






1-dim
←
iFFT






Sparse dimension-incremental FFT - method




Sparse dimension-incremental FFT - method







Sparse dimension-incremental FFT - method






construct
→
sampling set




Sparse dimension-incremental FFT - method






Sparse dimension-incremental FFT - method







1-dim
←
iFFT







Sparse dimension-incremental FFT - method




Sparse dimension-incremental FFT - method







Sparse dimension-incremental FFT - method







reconstructing
rank-1 lattice





Sparse dimension-incremental FFT - method






Sparse dimension-incremental FFT - method






Sparse dimension-incremental FFT - method




Sparse dimension-incremental FFT - method






Sparse dimension-incremental FFT - method




construct
→
sampling set






Sparse dimension-incremental FFT - method




1-dim
←
iFFT





Sparse dimension-incremental FFT - method






1-dim
←
iFFT






Sparse dimension-incremental FFT - method




Sparse dimension-incremental FFT - method






Sparse dimension-incremental FFT - method




reconstructing
rank-1 lattice






Sparse dimension-incremental FFT - method




1-dim
←
iFFT



Sparse dimension-incremental FFT - method



1-dim
←
iFFT



Sparse dimension-incremental FFT - results

search domain $\Gamma = \hat{G}_N^d$ full grid, $\sqrt{N} \lesssim |I| \lesssim N^d$

- samples: $\mathcal{O}(|I|^2 \log |\Gamma|)$
(1 iteration)
- computational costs: $\mathcal{O}(d|I|^3 + |I|^2(\log |\Gamma|) \log(|I| \log |\Gamma|))$
(1 iteration)
- for arbitrary Fourier coefficients $\hat{p}_{\mathbf{k}} \in \mathbb{C}$:
probabilistic approach with several iterations
- if ($\text{Re}(\hat{p}_{\mathbf{k}})$ identical sign) AND ($\text{Im}(\hat{p}_{\mathbf{k}})$ identical sign)
then deterministic version with 1 iteration

Sparse dimension-incremental FFT - example

- B-spline $N_m(x) := \sum_{k \in \mathbb{Z}} C_m \operatorname{sinc}\left(\frac{\pi}{m}k\right)^m \cos(\pi k) e^{2\pi i k x}$,
 $\|N_m\|_{L^2(\mathbb{T})} = 1$, $|\hat{N}_m(k)| \sim |k|^{-m}$
- $f(\mathbf{x}) := \prod_{t \in \{1,3,8\}} N_2(x_t) + \prod_{t \in \{2,5,6,10\}} N_4(x_t) + \prod_{t \in \{4,7,9\}} N_6(x_t)$

Sparse dimension-incremental FFT - example

- B-spline $N_m(x) := \sum_{k \in \mathbb{Z}} C_m \operatorname{sinc}\left(\frac{\pi}{m}k\right)^m \cos(\pi k) e^{2\pi i k x}$,
 $\|N_m\|_{L^2(\mathbb{T})} = 1$, $|\hat{N}_m(k)| \sim |k|^{-m}$
- $f(\mathbf{x}) := \prod_{t \in \{1,3,8\}} N_2(x_t) + \prod_{t \in \{2,5,6,10\}} N_4(x_t) + \prod_{t \in \{4,7,9\}} N_6(x_t)$
- full grid for $N = 64$, $d = 10$: $|\hat{G}_{64}^{10}| = 129^{10} \approx 1.28 \cdot 10^{21}$
- symmetric hyperbolic cross: $|I_{64}^{10}| = 696\,036\,321$
relative $L^2(\mathbb{T}^d)$ -error (best case) 4.1e-04

Sparse dimension-incremental FFT - example

- B-spline $N_m(x) := \sum_{k \in \mathbb{Z}} C_m \operatorname{sinc}\left(\frac{\pi}{m}k\right)^m \cos(\pi k) e^{2\pi i k x}$,
 $\|N_m\|_{L^2(\mathbb{T})} = 1$, $|\hat{N}_m(k)| \sim |k|^{-m}$
- $f(\mathbf{x}) := \prod_{t \in \{1,3,8\}} N_2(x_t) + \prod_{t \in \{2,5,6,10\}} N_4(x_t) + \prod_{t \in \{4,7,9\}} N_6(x_t)$
- full grid for $N = 64$, $d = 10$: $|\hat{G}_{64}^{10}| = 129^{10} \approx 1.28 \cdot 10^{21}$
- symmetric hyperbolic cross: $|I_{64}^{10}| = 696\,036\,321$
relative $L^2(\mathbb{T}^d)$ -error (best case) 4.1e-04
- results for dimension incremental algorithm with $\Gamma = \hat{G}_{64}^{10}$:

threshold	#samples	$ I $	rel. L_2 -error
1.0e-02	254 530	491	1.4e-01
1.0e-03	2 789 050	1 121	1.1e-02
1.0e-04	17 836 042	3 013	1.7e-03
1.0e-05	82 222 438	7 163	4.7e-04

Non-periodic case

results can be transferred from periodic to non-periodic case:

- multivariate algebraic polynomial $p : [-1, 1]^d \rightarrow \mathbb{R}$ in Chebyshev form with frequencies $\text{supp. on } I \subset \mathbb{N}_0^d, |I| < \infty,$

$$a(\mathbf{x}) = \sum_{\mathbf{k} \in I} \hat{a}_{\mathbf{k}} \prod_{t=1}^d T_{k_t}(x_t), \quad \hat{a}_{\mathbf{k}} \in \mathbb{R},$$

	periodic	non-periodic
basis function	$e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$	$\prod_{t=1}^d T_{k_t}(x_t)$

Non-periodic case

results can be transferred from periodic to non-periodic case:

- multivariate algebraic polynomial $p : [-1, 1]^d \rightarrow \mathbb{R}$ in Chebyshev form with frequencies supp. on $I \subset \mathbb{N}_0^d$, $|I| < \infty$,

$$a(\mathbf{x}) = \sum_{\mathbf{k} \in I} \hat{a}_{\mathbf{k}} \prod_{t=1}^d T_{k_t}(x_t), \quad \hat{a}_{\mathbf{k}} \in \mathbb{R},$$

	periodic	non-periodic
basis function	$e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$	$\prod_{t=1}^d T_{k_t}(x_t)$
spatial nodes	rank-1 lattice $\mathbf{x}_j = \frac{j}{M} \mathbf{z} \bmod \mathbf{1}$	rank-1 Chebyshev lattice $\mathbf{x}_j = \cos\left(\frac{j}{M} \pi \mathbf{z}\right)$ [Poppe, Cools 12]

Non-periodic case

results can be transferred from periodic to non-periodic case:

- multivariate algebraic polynomial $p : [-1, 1]^d \rightarrow \mathbb{R}$ in Chebyshev form with frequencies supp. on $I \subset \mathbb{N}_0^d$, $|I| < \infty$,

$$a(\mathbf{x}) = \sum_{\mathbf{k} \in I} \hat{a}_{\mathbf{k}} \prod_{t=1}^d T_{k_t}(x_t), \quad \hat{a}_{\mathbf{k}} \in \mathbb{R},$$

	periodic	non-periodic
basis function	$e^{2\pi i \mathbf{k} \cdot \mathbf{x}}$	$\prod_{t=1}^d T_{k_t}(x_t)$
spatial nodes	rank-1 lattice $\mathbf{x}_j = \frac{j}{M} \mathbf{z} \bmod \mathbf{1}$	rank-1 Chebyshev lattice $\mathbf{x}_j = \cos\left(\frac{j}{M} \pi \mathbf{z}\right)$ [Poppe, Cools 12]
evaluation / reconstruction	1-dim. FFT [Li, Hickernell 03] / [Kämmerer, Kunis, Potts 12]	1-dim. DCT [Cools, Poppe 11] [Potts, V. 15] / [Kämmerer 13] [Potts, V. 15]

Non-periodic case - related work

polynomial approximation on Lissajous curves [Bos, De Marchi, Vianello 15]

- $l_{\mathbf{z}}(t) := (\cos(z_1 t), \dots, \cos(z_d t)), t \in [0, \pi]$
- rank-1 Chebyshev lattice if $t = 0, \pi/M, 2\pi/M, \dots, \pi$
- results for algebraic polynomials of total or maximum degree $\leq n$

Non-periodic case - related work

polynomial approximation on Lissajous curves [Bos, De Marchi, Vianello 15]

- $\ell_{\mathbf{z}}(t) := (\cos(z_1 t), \dots, \cos(z_d t)), t \in [0, \pi]$
- rank-1 Chebyshev lattice if $t = 0, \pi/M, 2\pi/M, \dots, \pi$
- results for algebraic polynomials of total or maximum degree $\leq n$
- case $d = 2$
 - Padua point set,
e.g. [Bos, Caliari, Marchi, Vianello, Xu 06]
 $\mathcal{A}_n := \{\mathbf{x}_j := (\cos(j\pi/(n+1)), \cos(j\pi/n))^\top : j = 0, \dots, M\}$
 - $\mathcal{A}_n = \text{CL}(\mathbf{z}, M)$, where $\mathbf{z} := (n, n+1)^\top$ and $M := n(n+1)$
 - exact reconstruction of arbitrary 2d algebraic polynomial of total degree $\leq n$

Relation to tent-transformed rank-1 lattices


- tent-transformed rank-1 lattices

$$P_\psi(\mathbf{z}, M) := \{\psi(j\mathbf{z}/M \bmod \mathbf{1}) : j = 0, \dots, M-1\}$$


[Dick, Nuyens, Pillichshammer 14], [Suryanarayana, Nuyens, Cools 15],

[Cools, Kuo, Nuyens, Suryanarayana 16]


- tent transform $\psi: [0, 1] \rightarrow [0, 1]$, $\psi(x) := 1 - |2x - 1|$
- functions $g: [0, 1]^d \rightarrow \mathbb{R}$, $g(\mathbf{x}) := \sum_{\mathbf{k} \in I} g_{\mathbf{k}} \phi_{\mathbf{k}}(\mathbf{x})$, $I \subset \mathbb{N}_0^d$,
- half-periodic cosine functions $\phi_{\mathbf{k}}(\mathbf{x}) := \prod_{t=1}^d \sqrt{2}^{\delta_{0,k_t}} \cos(\pi k_t x_t)$,
 $\mathbf{k} \in \mathbb{N}_0^d$, ONB of $L_2([0, 1]^d)$
- reconstructing rank-1 lattice $\text{R1L}(\mathbf{z}, M)$ for $\mathcal{M}(I)$
- reconstruction from samples $g(P_\psi(\mathbf{z}, M))$ with 1-dim FFT
- polynomials g are not algebraic polynomials in general




I



$\mathcal{M}(I)$



reco. $\text{R1L}(\mathbf{z}, M)$



$P_\psi(\mathbf{z}, M)$

- **known** (arbitrary) frequency index set via rank-1 lattices
 - fast evaluation / reconstruction of **trigonometric polynomials**
[Li, Hickernell 03] / [Kämmerer, Kunis, Potts 12] [Kämmerer 13]
 - approximation of **periodic functions**
[Kuo, Sloan, Woźniakowski 06] [Kämmerer, Potts, V. 15] [Byrenheid, Kämmerer, Ullrich, V. 16]
 - fast evaluation / reconstruction of **algebraic polynomials**
[Cools, Poppe 11] [Potts, V. 15] / [Potts, V. 15]
 - approximation of **non-periodic functions**
[Dick, Nuyens, Pillichshammer 14] [Suryanarayana, Nuyens, Cools 15]
[Cools, Kuo, Nuyens, Suryanarayana 16]
- **unknown** frequency index set via rank-1 lattices
 - high-dim. sparse FFT via rank-1 lattice and 1-dim. sparse FFT
e.g. [Potts, Tasche, V. 16]
 - sparse dimension-incremental FFT based on rank-1 lattices
(periodic and non-periodic)
[Potts, V. 15] [V. 17]